WEBVTT

- 1.00:00:00.080 --> 00:00:02.010 < v -> Today it is my honor to introduce, <math></v>
- $2~00:00:02.010 \longrightarrow 00:00:04.309~Dr.$ Atul Deshpande.
- $3~00:00:04.309 \longrightarrow 00:00:06.790$ Dr. Deshpande is a postdoctoral researcher
- $4\ 00:00:06.790 \longrightarrow 00:00:09.080$ in the lab of Dr. Elana Fertig
- 5 00:00:09.080 --> 00:00:10.630 in the department of oncology,
- 6~00:00:10.630 -> 00:00:12.920 at Johns Hopkins University.
- 7 00:00:12.920 --> 00:00:14.730 He has a PhD in electrical engineering
- 8 00:00:14.730 --> 00:00:17.460 from the University of Wisconsin-Madison,
- $9\ 00:00:17.460 \longrightarrow 00:00:18.380$ and his interests include
- $10\ 00:00:18.380 \longrightarrow 00:00:20.100$ the use of time series analysis
- 11 00:00:20.100 --> 00:00:21.310 and spatial statistics
- $12\ 00:00:21.310 \longrightarrow 00:00:23.920$ for modeling biological processes.
- 13 00:00:23.920 --> 00:00:26.080 He's currently developing analysis techniques
- $14\ 00:00:26.080 \longrightarrow 00:00:28.027$ to use single cell and spacial multigenomics
- $15\ 00:00:28.027 \longrightarrow 00:00:30.110$ for the characterization of
- $16\ 00:00:30.110 \longrightarrow 00:00:31.509$ the tumor microenvironment
- $17\ 00:00:31.509 \longrightarrow 00:00:34.340$ and intracellular signaling networks.
- 18 00:00:34.340 --> 00:00:37.413 Welcome. (students applause)
- $19\ 00:00:40.090 --> 00:00:40.960 < v -> Well, thank you so much. </v>$
- $20\ 00:00:40.960 \longrightarrow 00:00:43.160$ And once I figure out my...
- 21 00:00:48.010 --> 00:00:49.380 Where my PowerPoint window is,
- $22\ 00:00:49.380 \longrightarrow 00:00:51.860$ we can start in earnest.
- $23~00{:}00{:}51.860 \dashrightarrow 00{:}00{:}54.940$ Okay, yeah, thank you for the kind introduction.
- $24\ 00:00:54.940 \longrightarrow 00:00:56.830$ So, I'm Atul Deshpande,
- $25\ 00:00:56.830 \longrightarrow 00:01:00.850$ and today the title of my talk is exploring time
- $26\ 00:01:00.850 \longrightarrow 00:01:03.640$ and space for identifying gene interactions
- 27 00:01:03.640 --> 00:01:05.340 using single cell transcriptomics.
- 28 00:01:06.630 --> 00:01:10.440 So, what do time and space mean
- 29 00:01:10.440 --> 00:01:12.649 in the context of this talk?

- $30\ 00:01:12.649 \longrightarrow 00:01:14.820$ So, they refer to recent technological advances
- 31 00:01:14.820 --> 00:01:17.070 and the algorithms, which are the foundation
- 32 00:01:17.070 --> 00:01:19.120 for the projects I will be talking about.
- $33\ 00:01:20.450 \longrightarrow 00:01:23.540$ And the first advance is the ability
- $34\ 00:01:23.540 \longrightarrow 00:01:26.730$ to measure gene expression in individual cells.
- $35\ 00:01:26.730 --> 00:01:28.790$ This in turn inspired development
- $36~00:01:28.790 \longrightarrow 00:01:31.740$ of algorithms that ordered these cells along
- $37\ 00:01:31.740 \longrightarrow 00:01:33.083$ the biological trajectory.
- $38\ 00:01:34.060 \longrightarrow 00:01:37.460$ Using these algorithms, we can observe changes
- 39 00:01:37.460 --> 00:01:39.440 in gene expression in
- 40 00:01:39.440 --> 00:01:42.850 a pseudo temporal reference for pseudo time,
- $41\ 00:01:42.850 \longrightarrow 00:01:44.870$ which is a measure of the progress
- $42\ 00:01:44.870 \longrightarrow 00:01:46.653$ of the biological process.
- 43 00:01:47.880 --> 00:01:50.170 The second is a more recent ability
- $44\ 00:01:50.170 \longrightarrow 00:01:51.500$ to measure gene expression
- $45\ 00:01:51.500 \longrightarrow 00:01:54.210$ within the spatial context of the tissue.
- $46\ 00:01:54.210 \longrightarrow 00:01:55.770$ But this we can analyze changes
- 47 00:01:55.770 --> 00:01:56.840 in gene expression
- 48 00:01:58.300 --> 00:02:00.450 as cellular neighborhoods change,
- $49\ 00:02:00.450 \longrightarrow 00:02:02.183$ or as the tissue type changes.
- $50\ 00:02:06.620 --> 00:02:09.800$ So, before single cell transcriptomics,
- $51\ 00:02:09.800 \longrightarrow 00:02:11.550$ we would usually get one measurement
- $52\ 00:02:11.550 \longrightarrow 00:02:15.300$ of gene expression from a collected sample.
- $53\ 00:02:15.300 \longrightarrow 00:02:18.300$ And this is now called
- 54 00:02:18.300 --> 00:02:22.450 bulk RNA-seq in retroactively.
- $55\ 00:02:22.450 \longrightarrow 00:02:25.510$ However, as this measurement would just be
- $56\ 00:02:25.510 --> 00:02:27.480$ an average of the population of cells
- $57\ 00:02:27.480 \longrightarrow 00:02:30.740$ in the sample, and it would obscure information
- $58\ 00:02:30.740 \longrightarrow 00:02:33.550$ about the different cell types, or different
- $59\ 00:02:33.550 \longrightarrow 00:02:35.093$ cell states in the population.
- 60 00:02:36.060 --> 00:02:37.490 With single-cell RNA-seq,

- $61\ 00:02:37.490 --> 00:02:39.510$ we can now measure gene expression
- $62\ 00:02:39.510 \longrightarrow 00:02:41.390$ in individual cells.
- 63 00:02:41.390 --> 00:02:43.180 Depending on technology, this can range
- $64\ 00:02:43.180 --> 00:02:46.320$ from a few hundred cells up to hundreds
- $65\ 00:02:46.320 \longrightarrow 00:02:48.690$ of thousands of cells.
- $66\ 00:02:48.690 \longrightarrow 00:02:51.000$ And this allows us to observe
- $67\ 00:02:51.000 \longrightarrow 00:02:54.680$ the full heterogeneity of the cell population
- $68\ 00:02:56.000 --> 00:02:58.510$ represented by gene expression.
- $69\ 00:02:58.510 \longrightarrow 00:03:01.770$ And using this high dimensional data
- $70\ 00:03:01.770 \longrightarrow 00:03:03.240$ that we now have,
- $71\ 00:03:03.240 \longrightarrow 00:03:05.320$ we can characterize different cell types
- $72\ 00:03:05.320 \longrightarrow 00:03:10.320$ and cell states as gene expression vectors.
- $73\ 00:03:11.670 --> 00:03:13.740$ So, one drawback of this technique
- $74~00:03:13.740 \longrightarrow 00:03:16.680$ is the issue of technical dropouts.
- 75 00:03:16.680 --> 00:03:20.790 Now, this is characterized by observing,
- $76\ 00:03:20.790 \longrightarrow 00:03:23.040$ as in us observing a lot
- 77 00:03:23.040 --> 00:03:26.300 of false zeroes, or zero inflated measurements,
- $78\ 00:03:26.300 --> 00:03:29.050$ because we are unable to reliably measure
- $79\ 00:03:29.050 \longrightarrow 00:03:31.213$ the low iron accounts in individual cells.
- $80\ 00:03:34.696 \longrightarrow 00:03:36.529$ Now, the first project
- 81 00:03:39.330 --> 00:03:41.520 that I will discuss uses
- 82 00:03:43.204 --> 00:03:46.430 a single cell RNA-seq technology,
- $83\ 00:03:46.430 \longrightarrow 00:03:48.800$ or as it's downstream of that.
- 84 00:03:48.800 --> 00:03:52.900 And it uses also downstream of algorithms,
- $85\ 00:03:52.900 \longrightarrow 00:03:57.900$ which order single cell data into trajectories,
- 86 00:03:57.970 --> 00:04:00.520 which represent the biology
- $87\ 00:04:00.520 \longrightarrow 00:04:02.220$ that they might be studying.
- 88 00:04:02.220 --> 00:04:04.280 For example, let's say if you are...
- 89 00:04:04.280 --> 00:04:08.580 You have a dataset, which corresponds
- $90\ 00:04:08.580 \longrightarrow 00:04:10.570$ to stem cell differentiation,
- $91\ 00:04:10.570 --> 00:04:13.480$ there are probably now 70 different

- 92 00:04:15.366 --> 00:04:17.030 trajectory inference methods depending on what
- 93 00:04:17.030 --> 00:04:20.900 kind of datasets you are studying,
- 94 00:04:20.900 --> 00:04:23.090 what biology you want to study,
- $95\ 00:04:23.090 \longrightarrow 00:04:24.920$ how big the dataset is,
- $96\ 00:04:24.920 \longrightarrow 00:04:27.520$ or what the expected trajectory is
- 97 00:04:27.520 --> 00:04:30.470 of the biology that you're studying maybe.
- $98~00:04:30.470 \longrightarrow 00:04:33.700$ And they attempt to order these cells based
- 99 00:04:33.700 --> 00:04:36.500 on the expression of potentially
- 100 00:04:36.500 --> 00:04:40.390 a few key marker genes, or how, which genes
- $101\ 00:04:40.390 \longrightarrow 00:04:42.870$ are differentially expressed along
- $102\ 00:04:42.870 \longrightarrow 00:04:44.363$ the biological process.
- 103 00:04:45.630 --> 00:04:47.620 So, anytime you collect,
- 104 00:04:47.620 --> 00:04:51.200 let's say a single cell RNA-seq data,
- 105 00:04:51.200 --> 00:04:54.460 you would find a mix of cells,
- $106\ 00:04:54.460 --> 00:04:55.900$ and that was the entire motivation
- $107\ 00:04:55.900 \longrightarrow 00:04:57.180$ for doing this.
- 108 00:04:57.180 --> 00:05:01.340 But that mix of cells would have
- $109\ 00:05:01.340 \longrightarrow 00:05:03.440$ a range of cell states,
- 110 00:05:03.440 --> 00:05:05.960 which could correspond to
- 111 00:05:06.860 --> 00:05:08.960 from the beginning of the biological process,
- $112\ 00:05:08.960 \longrightarrow 00:05:11.610$ to the very end of the biological process.
- $113\ 00:05:11.610 --> 00:05:14.480$ And what these algorithms are trying to do
- $114\ 00:05:14.480 \longrightarrow 00:05:18.050$ is they're trying to fit these cells
- $115\ 00:05:18.050 \longrightarrow 00:05:22.500$ in their right place, in the biological process.
- $116\ 00:05:22.500 \longrightarrow 00:05:25.490$ And once we do that, we can actually observe
- $117\ 00:05:25.490 \longrightarrow 00:05:30.250$ the gene expression along this ordering.
- $118\ 00:05:30.250 \longrightarrow 00:05:34.110$ And a lot of these methods also assign
- $119\ 00:05:34.110 \longrightarrow 00:05:35.450$ a pseudo time to each cell,
- $120\ 00:05:35.450 --> 00:05:38.810$ which tells you how far along in the biology

- 121 00:05:38.810 \rightarrow 00:05:43.460 they think, or they hypothesize that the cell is
- $122\ 00:05:43.460 \longrightarrow 00:05:44.660$ And so, the question that we wanted
- 123 00:05:44.660 --> 00:05:49.660 to ask is given this pseudo temporal ordering
- $124\ 00:05:50.290 \longrightarrow 00:05:52.660$ of the cells, which gives us
- $125\ 00:05:53.560 --> 00:05:55.140$ a gene expression dynamics
- $126\ 00:05:55.140 --> 00:05:57.970$ in the pseudo temporal reference.
- $127\ 00:05:57.970 \longrightarrow 00:06:01.760$ Can we use these dynamics
- $128\ 00:06:02.730 \longrightarrow 00:06:05.803$ to infer gene regulatory networks?
- 129 00:06:06.980 --> 00:06:10.100 Or any directed networks from say,
- $130\ 00:06:10.100 \longrightarrow 00:06:13.500$ sets of genes to their targets.
- $131\ 00:06:13.500 \longrightarrow 00:06:14.360$ And the second question
- $132\ 00{:}06{:}14.360 \dashrightarrow 00{:}06{:}19.250$ was whether the assigned pseudo time values help
- $133\ 00:06:19.250 \longrightarrow 00:06:22.313$ us in the network inference task.
- $134\ 00:06:25.370 \longrightarrow 00:06:30.370$ So, to make the, I guess,
- 135 00:06:31.470 --> 00:06:33.750 explanation more approachable,
- $136\ 00:06:33.750 \longrightarrow 00:06:36.523$ I will just use an example dataset.
- 137 00:06:37.380 --> 00:06:41.220 And as I explained, the concepts I've...
- $138\ 00:06:41.220 \longrightarrow 00:06:43.540$ We will just see what that means
- $139\ 00:06:43.540 \longrightarrow 00:06:45.400$ in terms of this dataset.
- 140 00:06:45.400 --> 00:06:49.650 So, this is a dataset from Semrau et al,
- $141\ 00:06:49.650 \longrightarrow 00:06:52.530$ and this is a single cell data
- 142 00:06:52.530 --> 00:06:56.730 from retinoic acid, driven differentiation.
- 143 00:06:56.730 --> 00:07:00.740 And in this mouse, embryonic stem cells
- $144\ 00:07:00.740 --> 00:07:02.456$ differentiate into neuroectoderm
- $145\ 00:07:02.456 --> 00:07:05.570$ and extraembryonic endoderm cells.
- 146 00:07:05.570 --> 00:07:09.790 Now the data as collected had nine samples,
- $147\ 00:07:09.790 \longrightarrow 00:07:12.090$ one before the differentiation starts
- $148\ 00:07:12.090 --> 00:07:15.180$ and one after every six hours.
- 149 00:07:15.180 --> 00:07:18.650 So, you have data collected over 96 hours

- $150\ 00:07:18.650 \dashrightarrow 00:07:22.453$ from nine samples, and each sample has 384 cells.
- 151 00:07:23.960 --> 00:07:26.530 So overall, I believe we have something
- $152\ 00:07:26.530 \longrightarrow 00:07:28.750$ like you can do the math.
- $153\ 00:07:28.750 \longrightarrow 00:07:31.793\ I guess,\ 2,600\ cells$ or something like that.
- $154\ 00:07:33.230 \longrightarrow 00:07:37.240$ So, we chose to apply
- $155\ 00:07:37.240 \longrightarrow 00:07:39.380$ two trajectory inference methods to this.
- $156\ 00:07:39.380 \longrightarrow 00:07:41.630$ So, the first one is monocle 2,
- $157\ 00:07:41.630 \longrightarrow 00:07:44.800$ which is also called Monocle DDR tree, I believe.
- 158 00:07:44.800 --> 00:07:46.960 And the second one is PAGA Tree.
- $159\ 00:07:46.960 \longrightarrow 00:07:50.400$ So, both of these methods identify
- $160\ 00:07:50.400 \longrightarrow 00:07:53.480$ a bifurcating trajectory from these cells.
- $161\ 00:07:53.480 \longrightarrow 00:07:56.170$ And so, the first one is to the left
- $162\ 00:07:56.170 --> 00:08:00.570$ where the embryonic stem cells are actually
- $163\ 00:08:00.570 \longrightarrow 00:08:01.910$ on the right of...
- $164\ 00{:}08{:}03.340 \dashrightarrow 00{:}08{:}07.400$ I'm not sure if people can see my mouse pointer,
- 165 00:08:07.400 --> 00:08:09.300 but yeah, they're on the right of the trajectory.
- 166 00:08:09.300 --> 00:08:13.520 And then, towards the bottom left,
- $167\ 00:08:13.520 --> 00:08:15.640$ you go into a neuroectoderm state
- $168\ 00:08:15.640 \longrightarrow 00:08:16.660$ and towards the...
- 169 00:08:18.770 --> 00:08:23.770 Right, top left, you go into an endoderm state.
- 170 00:08:24.390 --> 00:08:27.250 And on the right side, the way PAGA Tree
- 171 00:08:27.250 --> 00:08:29.880 infers trajectory is you have
- $172\ 00:08:29.880 \longrightarrow 00:08:32.980$ the embryonic stem cells on the top left.
- $173\ 00:08:32.980 \longrightarrow 00:08:35.100$ And then, it identifies
- $174\ 00:08:35.100 \longrightarrow 00:08:39.100$ a few more branches than Monocle does.
- 175 00:08:39.100 --> 00:08:40.230 But both of these
- 176 00:08:40.230 --> 00:08:43.030 identify branching trajectories.
- $177\ 00:08:43.030 \longrightarrow 00:08:47.000$ And in each case we selected
- 178 00:08:48.150 --> 00:08:49.210 the two branches,

- 179 00:08:49.210 --> 00:08:53.530 which corresponded to markers, which were,
- $180\ 00:08:53.530 --> 00:08:55.920$ which ended up being high for neuroectoderm.
- 181 00:08:55.920 --> 00:08:59.670 So, the trajectories, the sub trajectories
- 182 00:08:59.670 --> 00:09:02.810 from each method that we've wanted to study
- $183\ 00:09:02.810 \longrightarrow 00:09:07.420$ was the embryonic stem cells to neuroectoderm,
- $184\ 00:09:08.480 \longrightarrow 00:09:10.293$ using these two methods.
- $185\ 00:09:11.370 \longrightarrow 00:09:13.540$ So, this as in, so we had...
- $186\ 00:09:13.540 \longrightarrow 00:09:16.240$ We have these two trajectory inference methods,
- 187 00:09:16.240 --> 00:09:18.380 which assigned their own pseudo times,
- $188\ 00:09:18.380 \longrightarrow 00:09:22.500$ and this is the pseudo temporal expression
- $189\ 00:09:22.500 \longrightarrow 00:09:25.410$ dynamics for the same gene.
- 190 00:09:25.410 --> 00:09:28.670 I did not mark which gene it was, but yeah,
- $191\ 00:09:28.670 \longrightarrow 00:09:30.220$ so this was for the same gene.
- 192 00:09:30.220 --> 00:09:32.760 And you can see that the dynamics
- 193 00:09:32.760 --> 00:09:35.290 that each of these trajectories gives
- $194\ 00:09:35.290 \longrightarrow 00:09:36.680$ us is different.
- 195 00:09:36.680 --> 00:09:39.690 First of all, the main branch,
- 196 00:09:39.690 --> 00:09:41.820 or sub part of the trajectory that
- $197\ 00:09:41.820 --> 00:09:43.960$ we are considering has
- $198\ 00:09:43.960 \longrightarrow 00:09:45.610$ a different number of cells.
- $199\ 00{:}09{:}45.610 \dashrightarrow 00{:}09{:}47.940$ And these cells may not necessarily be common
- $200\ 00:09:47.940 \longrightarrow 00:09:48.800$ to both end.
- $201\ 00:09:48.800 --> 00:09:49.810$ There will be some which are common
- 202 00:09:49.810 --> 00:09:51.590 to both of these trajectories,
- $203\ 00{:}09{:}51.590 \dashrightarrow 00{:}09{:}54.240$ but some others which are completely different.
- 204 00:09:54.240 --> 00:09:56.580 But also, that the cell ordering itself
- $205\ 00{:}09{:}56.580 {\: \mbox{--}}{\:>}\ 00{:}10{:}01.320$ that each method based on whatever mathematics

```
206\ 00:10:01.320 \longrightarrow 00:10:03.420 they use, or whatever algorithms they use,
```

- $207\ 00:10:04.600 \longrightarrow 00:10:07.780$ would differ between these two methods.
- $208\ 00:10:07.780 --> 00:10:11.840$ So, as you see, Monocle has a higher expression
- 209 00:10:11.840 --> 00:10:13.620 much earlier in the pseudo time,
- $210~00{:}10{:}13.620 \dashrightarrow 00{:}10{:}17.820$ as opposed to PAGA Tree, which has much later.
- 211 00:10:17.820 --> 00:10:20.020 And the pseudo times here,
- 212 00:10:20.020 --> 00:10:22.010 were not exactly 100, they're just nominalized
- 213 00:10:22.010 --> 00:10:25.470 to 100 just represent progress from 0%
- 214 00:10:25.470 --> 00:10:27.863 of the biology to 100% of the biology,
- $215\ 00:10:30.620 \longrightarrow 00:10:32.973$ or as inferred by that method.
- 216 00:10:33.810 --> 00:10:36.980 So, now what are the challenges associated
- $217\ 00:10:36.980 \longrightarrow 00:10:39.400$ with order single-cell data?
- 218 00:10:39.400 --> 00:10:42.840 So, the first one is that unlike say,
- 219 00:10:42.840 --> 00:10:46.580 stock data, or say weather data,
- 220 00:10:46.580 --> 00:10:49.130 or something like that, you don't necessarily
- $221\ 00:10:49.130 \longrightarrow 00:10:53.630$ have a uniform distribution of cells.
- $222\ 00:10:53.630 --> 00:10:56.330$ And if you're going to do a time series analysis,
- 223 00:10:56.330 --> 00:10:57.480 that would mean that you do not
- 224 00:10:57.480 --> 00:10:59.820 have regularly spaced time series,
- 225 00:10:59.820 --> 00:11:00.653 but you actually
- 226 00:11:00.653 --> 00:11:03.020 have irregularly space time series.
- 227 00:11:03.020 --> 00:11:05.090 On top of that, the pseudo time values
- $228\ 00:11:05.090 \longrightarrow 00:11:07.270$ that are assigned to the cells
- $229~00{:}11{:}07.270 \dashrightarrow 00{:}11{:}10.133$ and ordering stem cells is uncertain.
- 230 00:11:12.720 --> 00:11:16.720 Now, finally, we recall that we had the issue
- 231 00:11:16.720 --> 00:11:18.740 of zero inflated measurements,
- $232\ 00:11:18.740 \longrightarrow 00:11:20.870$ or false zeroes in the meter
- $233\ 00:11:20.870 \longrightarrow 00:11:22.370$ because of technical dropouts.
- $234\ 00:11:25.798 \longrightarrow 00:11:28.720$ So, the question is how to overcome all
- $235\ 00:11:28.720 \longrightarrow 00:11:32.370$ of these drawbacks

```
236 00:11:32.370 --> 00:11:33.720 to try and find
```

- $237\ 00:11:35.730 --> 00:11:39.293$ networks from this time series data.
- 238 00:11:40.160 --> 00:11:43.310 So, the project that we had,
- $239\ 00:11:43.310 \longrightarrow 00:11:45.000$ it resulted in basically
- 240 00:11:45.000 --> 00:11:46.270 an algorithm called SINGE,
- $241\ 00:11:46.270 \longrightarrow 00:11:48.050$ which is single cell inference
- $242\ 00:11:48.050 \longrightarrow 00:11:50.210$ of networks from Granger ensembles.
- 243 00:11:50.210 --> 00:11:52.730 So, this was done at the Morgridge Institute
- 244 00:11:52.730 --> 00:11:55.160 for Research in Madison, Wisconsin.
- $245\ 00:11:55.160 \longrightarrow 00:11:58.633$ And these are my collaborators on this project.
- 246 00:12:00.770 --> 00:12:02.860 And let's see, okay.
- 247 00:12:02.860 --> 00:12:06.390 So, the main concept that we build on
- $248\ 00:12:06.390 \longrightarrow 00:12:08.400$ is basically the Granger causality test.
- $249\ 00:12:08.400 \longrightarrow 00:12:11.603$ It was introduced by Clive Granger in 1960s.
- 250 00:12:14.489 --> 00:12:15.550 And to give a very simple example
- 251 00:12:15.550 --> 00:12:17.330 of what it's trying to say is, let's say
- 252 00:12:17.330 --> 00:12:21.670 if you have two times series X and Y,
- 253 00:12:21.670 --> 00:12:23.970 now Granger causality tests, whether
- 254 00:12:25.940 --> 00:12:28.330 the prediction of current values of Y
- 255 00:12:28.330 --> 00:12:30.860 improves by using past values of X,
- $256\ 00:12:30.860 \longrightarrow 00:12:32.703$ in addition to past values of Y.
- 257 00:12:34.210 --> 00:12:35.870 And if that happens, then we say
- $258\ 00:12:35.870 \longrightarrow 00:12:37.940$ that X Granger causes Y.
- $259\ 00:12:37.940 \longrightarrow 00:12:40.590$ So, this is basically a lag regression
- $260\ 00:12:40.590 \longrightarrow 00:12:41.890$ between X and Y.
- 261 00:12:41.890 --> 00:12:43.860 So, this has had applications
- 262 00:12:43.860 --> 00:12:46.060 in econometrics and finance,
- $263\ 00:12:46.060 \longrightarrow 00:12:47.170$ and is also being used
- 264 00:12:47.170 --> 00:12:50.600 in computational neuroscience and biology,
- $265\ 00:12:50.600 \longrightarrow 00:12:53.643$ as noted in these examples here.
- $266\ 00:12:55.310 \longrightarrow 00:12:57.650$ Now, the multivariate Granger causality test

```
267 00:12:57.650 --> 00:13:00.290 can be thought of as setting up and solving
```

- $268\ 00:13:00.290 \longrightarrow 00:13:02.320$ a vector, or regression model,
- 269 00:13:02.320 --> 00:13:05.430 where you have say, P genes, T time points
- $270\ 00:13:05.430 \longrightarrow 00:13:06.440$ and L lags.
- 271 00:13:06.440 --> 00:13:09.073 Where L lags is telling you how many,
- $272\ 00{:}13{:}11.010 \dashrightarrow 00{:}13{:}13.760$ say your relationships with the past expressions
- $273\ 00:13:13.760 \longrightarrow 00:13:15.670$ you're trying to model.
- 274 00:13:15.670 --> 00:13:17.150 And once you have that,
- 275 00:13:17.150 --> 00:13:21.200 you could think of solving this way,
- 276 00:13:21.200 --> 00:13:23.740 our model by just minimizing
- $277\ 00:13:23.740 \longrightarrow 00:13:25.253$ this objective function here.
- 278 00:13:26.610 --> 00:13:28.060 And that would give you, I guess,
- $279\ 00:13:28.060 --> 00:13:30.700$ a few edges between the past values
- $280\ 00:13:30.700 \longrightarrow 00:13:34.160$ of all of the genes and your target gene.
- 281 00:13:34.160 --> 00:13:35.610 Okay, maybe I should have explained
- $282\ 00:13:35.610 \longrightarrow 00:13:36.920$ this figure first.
- 283 00:13:36.920 --> 00:13:39.250 So, you have all the regular,
- $284\ 00:13:39.250 \longrightarrow 00:13:42.050$ all the possible regulators of a gene,
- $285\ 00:13:42.050 \longrightarrow 00:13:43.340$ and then you have a target gene,
- 286 00:13:43.340 --> 00:13:44.790 and you're trying to identify
- 287 00:13:46.470 --> 00:13:48.840 what explains what past values
- $288\ 00:13:48.840 \longrightarrow 00:13:51.400$ of any of these genes explains
- $289\ 00:13:51.400 \longrightarrow 00:13:53.263$ the current values of the target gene.
- $290\ 00:13:54.580 \longrightarrow 00:13:58.730$ And if you wanted to have
- 291 00:13:58.730 --> 00:14:01.710 a sparse representation of this network,
- $292\ 00:14:01.710 \longrightarrow 00:14:03.300$ or have an...
- 293 00:14:03.300 --> 00:14:05.030 Count only a few of the edges,
- 294 00:14:05.030 --> 00:14:08.210 you would introduce this by CT parameter,
- 295 00:14:08.210 --> 00:14:11.960 which would ensure that the edges from say,
- $296\ 00:14:11.960 \longrightarrow 00:14:14.590$ all of these genes to your target

- $297\ 00:14:14.590 \longrightarrow 00:14:15.840$ are not numerous.
- $298~00{:}14{:}15.840 \dashrightarrow 00{:}14{:}18.453$ And you can explain the biology in a few edges.
- 299 00:14:22.378 --> 00:14:25.543 Now, to counter the irregularity
- $300\ 00:14:26.590 \longrightarrow 00:14:29.480$ of the time series, we use
- $301\ 00:14:30.340 \longrightarrow 00:14:33.220$ an idea called Generalized Lasso Granger.
- 302 00:14:33.220 --> 00:14:36.452 So, what this does is,
- $303\ 00:14:36.452 \longrightarrow 00:14:39.110$ I'm not sure, maybe I have...
- 304 00:14:39.110 --> 00:14:43.530 Yeah, okay, so just to recall, right?
- 305 00:14:43.530 --> 00:14:45.630 So, you have a pseudo temporal data,
- $306\ 00:14:45.630 \longrightarrow 00:14:48.140$ which has irregular time series,
- $307\ 00:14:48.140 \longrightarrow 00:14:50.680$ and you have missing values,
- $308\ 00:14:50.680 \longrightarrow 00:14:54.150$ which show up as zeros here, right?
- $309\ 00:14:54.150 --> 00:14:59.150$ So, we want to adapt the Lasso Granger test
- $310\ 00:15:00.280 \longrightarrow 00:15:01.990$ for irregular time series.
- 311 00:15:01.990 --> 00:15:04.740 So, what was previously,
- 312 00:15:04.740 --> 00:15:07.320 basically coefficients from older samples
- $313\ 00:15:07.320 \longrightarrow 00:15:08.820$ in regular time series,
- $314\ 00:15:08.820 \longrightarrow 00:15:13.720$ now becomes coefficients from just timestamps
- $315\ 00:15:14.880 \longrightarrow 00:15:16.010$ in the past.
- $316\ 00:15:16.010 --> 00:15:17.730$ Because you might not necessarily have
- $317\ 00:15:17.730 \longrightarrow 00:15:19.553$ a sample at that point.
- 318 00:15:20.710 --> 00:15:25.357 Furthermore, we can rethink basically,
- $319\ 00:15:28.091 \longrightarrow 00:15:32.780$ the object to function as originally,
- $320\ 00:15:32.780 --> 00:15:34.890$ if it was a dot predict between
- $321\ 00:15:34.890 \longrightarrow 00:15:37.650$ the coefficients and the values
- $322\ 00:15:37.650 \longrightarrow 00:15:40.560$ of the gene expression,
- 323 00:15:40.560 --> 00:15:45.200 we rethink that as a weighted dot predict,
- $324\ 00:15:45.200 \longrightarrow 00:15:46.650$ where basically we...
- $325\ 00:15:47.590 \longrightarrow 00:15:48.840$ And this is the description
- $326\ 00:15:48.840 \longrightarrow 00:15:51.450$ of the weighted dot predict, where you use

- $327~00{:}15{:}51.450 {\: --> \:} 00{:}15{:}55.720$ a Gaussian kernel to weight the inputs
- $328~00:15:55.720 \longrightarrow 00:15:58.540$ pseudo product based on their proximity
- $329\ 00:15:58.540 \longrightarrow 00:16:01.830$ to the timestamps that you...
- $330\ 00:16:01.830 \longrightarrow 00:16:04.260$ That correspond to these coefficients.
- $331\ 00:16:04.260 \longrightarrow 00:16:07.740$ So, these ellipses here show kernels,
- $332\ 00:16:07.740 \longrightarrow 00:16:09.600\ I$ guess, they represent kernels.
- $333\ 00:16:09.600 \longrightarrow 00:16:12.310$ They don't necessarily stop at these bandwidths,
- $334\ 00:16:12.310 \longrightarrow 00:16:13.210$ but they just keep going
- 335 00:16:13.210 --> 00:16:15.740 because they're ghosting kernels.
- 336 00:16:15.740 --> 00:16:17.990 But these just represent the kernels,
- 337 00:16:17.990 --> 00:16:20.260 where basically, if you have
- 338 00:16:20.260 --> 00:16:22.490 a timestamp corresponding to coefficient
- 339 00:16:22.490 --> 00:16:25.480 and you have no sample at that timestamp,
- $340\ 00:16:25.480 \longrightarrow 00:16:26.440$ that doesn't necessarily mean
- $341\ 00:16:26.440 \longrightarrow 00:16:30.880$ that the input to the gene predict it is zero.
- $342\ 00:16:30.880 \longrightarrow 00:16:32.960$ So, basically what you would do is
- 343 00:16:32.960 --> 00:16:36.010 you would just look at a bin around
- $344\ 00:16:36.010 \longrightarrow 00:16:41.010$ that timestamp, and weight input from regulators,
- $345\ 00{:}16{:}42.180 \dashrightarrow 00{:}16{:}46.240$ depending on their proximity to this timestamp.
- $346\ 00:16:46.240 \longrightarrow 00:16:50.790$ So, if the sample is exactly at
- 347 00:16:50.790 --> 00:16:52.110 the timestamp that you expect,
- $348\ 00:16:52.110 \longrightarrow 00:16:54.400$ you would rate it highly based
- $349\ 00:16:54.400 \longrightarrow 00:16:56.440$ on discussion kernel, and the farther
- $350\ 00:16:56.440 --> 00:16:58.350$ you move away from the timestamp,
- $351\ 00:16:58.350 \longrightarrow 00:17:01.170$ the weaker the rate of
- $352\ 00:17:02.240 \longrightarrow 00:17:05.360$ that particular sample would be.
- $353\ 00:17:05.360 \longrightarrow 00:17:06.900$ So, what this helps us do
- $354\ 00:17:06.900 \longrightarrow 00:17:10.160$ is if there are say more than one cells
- 355 00:17:10.160 --> 00:17:13.870 in close proximity, it would take input

- $356\ 00:17:13.870 \longrightarrow 00:17:15.400$ from all of them.
- $357\ 00:17:15.400 \longrightarrow 00:17:17.670$ If there are no cells in the close proximity
- 358 00:17:17.670 --> 00:17:19.680 to at least take input from some cells,
- $359\ 00:17:19.680 \longrightarrow 00:17:21.380$ which are farther away, and so on.
- $360\ 00:17:24.510 \longrightarrow 00:17:27.340$ So, yeah, as in this works
- $361\ 00:17:27.340 \longrightarrow 00:17:28.460$ with irregular time series,
- $362\ 00:17:28.460 --> 00:17:30.370$ because you don't necessarily have
- $363\ 00{:}17{:}30.370 \dashrightarrow 00{:}17{:}33.500$ to expect samples in the past at the timestamps
- $364\ 00:17:33.500 \longrightarrow 00:17:34.700$ that you wanted them to.
- $365\ 00:17:36.480 --> 00:17:39.900$ And yeah, I think we already discussed this.
- $366~00{:}17{:}39.900 \dashrightarrow 00{:}17{:}44.900$ So, now, as in going back to the case for...
- $367\ 00:17:45.420 \longrightarrow 00:17:48.210$ So, we had these false zeroes, right?
- 368 00:17:48.210 --> 00:17:50.310 So now, because of this kernel method,
- $369\ 00:17:50.310 \longrightarrow 00:17:54.010$ we have an inherent imputation over missing data.
- 370 00:17:54.010 --> 00:17:56.423 So, now we get what we could think of as,
- 371 00:17:57.930 --> 00:18:00.400 instead of taking all of the zeros
- $372\ 00:18:00.400 \longrightarrow 00:18:02.550$ as they are at face value,
- $373\ 00:18:02.550 \longrightarrow 00:18:04.290$ we can treat them, or some of them
- $374\ 00:18:04.290 \longrightarrow 00:18:09.260$ as dropouts, as just missing data.
- 375 00:18:09.260 --> 00:18:11.230 And we just remove those samples now,
- $376\ 00:18:11.230 \longrightarrow 00:18:12.640$ because we can now work
- $377\ 00:18:12.640 \longrightarrow 00:18:14.820$ with irregular time series.
- 378 00:18:14.820 --> 00:18:17.170 And because of this kernel method,
- $379\ 00:18:17.170 \longrightarrow 00:18:19.420$ we can actually work with time signature,
- $380\ 00:18:19.420 --> 00:18:21.570$ all uniquely irregular.
- $381\ 00:18:21.570 \longrightarrow 00:18:22.890$ We can work with...
- $382\ 00:18:24.420 \longrightarrow 00:18:26.270$ We can remove the zero valued samples
- $383~00:18:26.270 \longrightarrow 00:18:29.860$ and get a different, differently irregular
- $384\ 00:18:29.860 \longrightarrow 00:18:31.853$ time series for each of these genes.

- $385\ 00:18:32.790 --> 00:18:36.870$ And so, such an action can probably
- $386~00{:}18{:}36.870 \dashrightarrow 00{:}18{:}39.750$ be informed by imputation techniques like magic,
- 387 00:18:39.750 --> 00:18:41.910 which help you complete,
- $388\ 00:18:41.910 \longrightarrow 00:18:43.730$ or impute zeros in the dataset.
- 389 00:18:43.730 --> 00:18:45.820 So, instead of imputing the dataset,
- $390\ 00:18:45.820 \longrightarrow 00:18:47.780$ as you could just use its output
- $391\ 00:18:47.780 --> 00:18:51.110$ to decide whether or not to remove the data from,
- $392\ 00:18:51.110 \longrightarrow 00:18:55.943$ or remove that zero from this input dataset.
- $393\ 00:18:58.140 \longrightarrow 00:18:59.930$ So, this is just an illustration
- $394\ 00:18:59.930 \longrightarrow 00:19:04.330$ of a single generalized Lasso Granger test.
- $395~00:19:04.330 \dashrightarrow 00:19:08.300$ So, you have the POU5F1 gene, and it's basically,
- $396\ 00:19:08.300 \longrightarrow 00:19:11.030$ you see it's the cells corresponding
- $397\ 00:19:11.030 \longrightarrow 00:19:15.860$ to that, or other details expression
- 398~00:19:15.860 --> 00:19:17.770 along pseudo time.
- 399 00:19:17.770 --> 00:19:22.010 And what you also see is two trendlines
- 400 00:19:22.950 --> 00:19:27.000 predicted using a Lambda of 0.1,
- $401\ 00:19:27.000 \longrightarrow 00:19:29.110$ which is basically a sparsity constraint of 0.1.
- 402 00:19:29.110 --> 00:19:31.990 So, it would have fewer edges
- $403\ 00:19:31.990 \longrightarrow 00:19:34.867$ between the regulators and POU5F1.
- $404\ 00:19:35.940 \longrightarrow 00:19:40.530$ And then a Lambda of 0.02,
- $405\ 00:19:40.530 \longrightarrow 00:19:42.640$ which has far more regulators.
- $406\ 00:19:42.640 \longrightarrow 00:19:45.570$ And you can see that both of these predict
- $407\ 00:19:46.460 \longrightarrow 00:19:49.350$ the trends of POU5F1 when using
- $408\ 00:19:49.350 \longrightarrow 00:19:50.943$ the past values quite well.
- $409\ 00:19:53.970 \longrightarrow 00:19:58.093$ So, now that was just one GLG test.
- $410\ 00{:}19{:}59.120 \dashrightarrow 00{:}20{:}01.460$ Now, what SINGE does, is it performs multiple
- $411\ 00:20:01.460 --> 00:20:04.310$ such GLG tests where you sub-sample
- $412\ 00:20:04.310 \longrightarrow 00:20:06.840$ the time series different ways

- $413\ 00:20:06.840 \longrightarrow 00:20:11.610$ to get different irregulars time series again.
- $414\ 00:20:11.610 \longrightarrow 00:20:14.110$ And you also use diverse hyper-parameters
- 415 00:20:14.110 --> 00:20:16.990 to effectively using these two combinations,
- 416 00:20:16.990 --> 00:20:20.220 slice the cake multiple ways and trying
- $417\ 00:20:20.220 \longrightarrow 00:20:21.720$ to look at the data.
- $418\ 00:20:21.720 \longrightarrow 00:20:22.890$ So, the type of barometers
- $419\ 00:20:22.890 \longrightarrow 00:20:25.210$ that we use are Lambda, which determines
- $420\ 00:20:25.210 \longrightarrow 00:20:28.650$ the sparsity of the network that we get,
- $421\ 00:20:28.650 \longrightarrow 00:20:30.830$ or get into metrics that we get.
- 422 00:20:30.830 --> 00:20:35.830 And we have Delta T, which gives us
- $423\ 00:20:35.900 \longrightarrow 00:20:39.530$ a time resolution of the lags between say,
- $424\ 00:20:39.530 \longrightarrow 00:20:41.280$ the past regulators
- 425 00:20:41.280 --> 00:20:44.990 and the current target timestamps,
- $426\ 00:20:44.990 \longrightarrow 00:20:47.360$ and the number of likes that you have.
- 427 00:20:47.360 --> 00:20:51.170 So together, they will tell you how far behind
- 428 00:20:51.170 --> 00:20:53.550 in pseudo time should you be looking to try
- $429\ 00:20:53.550 \longrightarrow 00:20:57.400$ to predict the expression of the target.
- 430 00:20:57.400 --> 00:20:58.680 And finally, the kernel width,
- $431\ 00:20:58.680 --> 00:21:03.030$ which tells how far, how wide the width should be
- $432\ 00{:}21{:}03.030 \dashrightarrow 00{:}21{:}06.813$ around the timestamp that you are considering.
- 433 00:21:08.456 --> 00:21:09.289 Now, once we get
- 434 00:21:11.370 --> 00:21:13.340 adjacency matrices from all of these,
- $435\ 00:21:13.340 \longrightarrow 00:21:16.860$ we get, we considered them as partial networks,
- $436\ 00:21:16.860 \longrightarrow 00:21:20.210$ and we get ranked lists from each of them.
- 437 00:21:20.210 --> 00:21:22.020 And we aggregate these rank lists
- $438\ 00:21:22.020 \longrightarrow 00:21:24.330$ using a modified border count.
- $439\ 00:21:24.330 \longrightarrow 00:21:25.390$ So, border count is something
- $440\ 00:21:25.390 \longrightarrow 00:21:28.806$ which has been used in election.
- 441 00:21:28.806 --> 00:21:31.420 It's basically an election, I guess,

```
442 00:21:31.420 --> 00:21:33.630 result aggregating strategy,
```

- 443 00:21:33.630 --> 00:21:35.950 where if you have five candidates,
- 444 00:21:35.950 --> 00:21:39.190 you rank them from one to five,
- 445 00:21:39.190 --> 00:21:41.600 and then the person who has, I guess,
- $446\ 00:21:41.600 \longrightarrow 00:21:44.360$ the lowest number here over all
- $447\ 00:21:44.360 \longrightarrow 00:21:46.630$ of the people that voted,
- $448\ 00:21:46.630 \longrightarrow 00:21:48.930$ they would win the vote.
- 449 00:21:48.930 --> 00:21:51.507 So, the modified border width
- $450\ 00:21:51.507 --> 00:21:53.190$ is basically the same concept,
- $451\ 00:21:53.190 \longrightarrow 00:21:55.260$ but the only change that we did
- $452\ 00:21:55.260 --> 00:22:00.260$ was we wanted to place more weight
- 453 00:22:02.870 --> 00:22:05.180 to a ranking, which distinguishes
- $454\ 00:22:05.180 \longrightarrow 00:22:09.580$ between say a one, the first interaction
- $455\ 00:22:09.580 \longrightarrow 00:22:12.310$ we find with the 10th interaction we find.
- $456\ 00:22:12.310 \longrightarrow 00:22:15.160$ As opposed to say, the 10,000th interaction
- $457\ 00:22:15.160 \longrightarrow 00:22:17.620$ we find with the 10,010th interaction
- $458\ 00:22:17.620 \longrightarrow 00:22:18.620$ that we find.
- 459 00:22:18.620 --> 00:22:23.320 So, that's why the weighting before adding
- 460 00:22:23.320 --> 00:22:26.210 these border weights is one over N squared,
- $461\ 00:22:26.210 \longrightarrow 00:22:29.243$ as opposed to say, N here.
- 462 00:22:33.100 --> 00:22:36.460 So, yeah, once we aggregate this,
- $463\ 00:22:36.460 \longrightarrow 00:22:39.310$ we get a final rank list.
- 464 00:22:39.310 --> 00:22:43.200 And so, we had to do in for trajectories,
- $465\ 00:22:43.200 \longrightarrow 00:22:45.770$ we got gene dynamics from them,
- $466\ 00:22:45.770 \longrightarrow 00:22:49.070$ and now that results in two different networks.
- $467\ 00:22:49.070 \longrightarrow 00:22:52.840$ And there's just showing the top 100 edges
- $468\ 00:22:52.840 \longrightarrow 00:22:54.910$ from Monocle 2 and PAGA Tree.
- 469 00:22:54.910 --> 00:22:56.320 Now, you can obviously see
- $470\ 00:22:56.320 \longrightarrow 00:22:58.600$ that they look very different.
- 471 00:22:58.600 --> 00:23:01.690 Some of the edges I think, are common,
- $472\ 00:23:01.690 \longrightarrow 00:23:04.660$ but they can be very, very different.

- $473\ 00:23:04.660 \longrightarrow 00:23:07.710$ So, now the question is,
- $474\ 00:23:07.710 \longrightarrow 00:23:11.520$ which of these is right, or better?
- $475\ 00:23:11.520 \longrightarrow 00:23:14.070$ So, for that we would have
- 476 00:23:14.070 --> 00:23:15.110 to first think of, okay,
- 477 00:23:15.110 --> 00:23:16.940 how do we evaluate this?
- $478\ 00:23:16.940 \longrightarrow 00:23:19.940$ So, one way to evaluate that would be
- $479\ 00:23:19.940 \longrightarrow 00:23:23.570$ to do a precision recall evaluation.
- 480 00:23:23.570 --> 00:23:25.370 So, let's say we have this rank list
- 481 00:23:25.370 --> 00:23:27.790 of candidate gene interactions that we just got
- 482 00:23:27.790 --> 00:23:30.680 from SINGE and a gold standard,
- $483\ 00:23:30.680 \longrightarrow 00:23:31.780$ which knows the truth.
- $484\ 00:23:32.680 \longrightarrow 00:23:34.390$ As we go down this rank list,
- $485\ 00:23:34.390 \longrightarrow 00:23:36.870$ the precision metric tells us
- $486\ 00:23:36.870 \longrightarrow 00:23:38.300$ what fraction of the prediction
- $487\ 00:23:38.300 \longrightarrow 00:23:40.370$ so far have been correct.
- $488\ 00:23:40.370 --> 00:23:41.710$ And the recall metric tells us
- $489\ 00:23:41.710 \longrightarrow 00:23:44.320$ how many of the total interactions
- 490 00:23:44.320 --> 00:23:46.110 in the gold standard, which were correct
- $491\ 00:23:46.110 \longrightarrow 00:23:47.433$ have so far been covered.
- 492 00:23:48.350 --> 00:23:50.880 So, the figure on the right shows
- 493 00:23:50.880 --> 00:23:53.940 a precision recall curve for two rank lists.
- $494\ 00:23:53.940 \longrightarrow 00:23:56.240$ The ideal precision recall curve
- $495\ 00:23:56.240 --> 00:23:58.220$ would place all the edges in the gold standard
- $496\ 00:23:58.220 \longrightarrow 00:23:59.053$ at the top of the list.
- 497 00:23:59.053 --> 00:24:03.560 So, that's the dotted line that you see here,
- $498\ 00:24:03.560 \longrightarrow 00:24:06.040$ and the area under that precision
- 499 00:24:06.040 --> 00:24:08.530 we call curve (mumbles) blue one.
- 500 00:24:08.530 --> 00:24:12.970 A random list in expectation would be flat.
- 501 00:24:12.970 --> 00:24:14.940 So, and it would have a precision
- 502 00:24:14.940 --> 00:24:17.930 recall curve, and the area under

- $503\ 00:24:17.930 \longrightarrow 00:24:20.260$ that curve would be 0.5.
- 504 00:24:20.260 --> 00:24:24.277 and here, I guess, to make belief orderings.
- $505\ 00:24:27.090 \longrightarrow 00:24:29.310$ And in this example, we can see
- 506 00:24:29.310 --> 00:24:34.063 that the precision we call curve of A,
- 507 00:24:35.150 --> 00:24:39.750 which I guess, the predictor A is better
- 508 00:24:39.750 --> 00:24:44.750 because it starts off with having more ones,
- $509\ 00:24:44.880 --> 00:24:47.930$ or as in a high precision, and then falls
- 510 00:24:47.930 --> 00:24:49.990 as opposed to B, which rises
- $511\ 00:24:49.990 --> 00:24:51.053$ from a low precision.
- $512\ 00:24:51.053 --> 00:24:54.600$ What it means that A gets more hits
- $513\ 00:24:54.600 \longrightarrow 00:24:56.860$ in the top of its list as opposed to B,
- $514\ 00:24:56.860 \longrightarrow 00:24:58.030$ and so on.
- $515\ 00:24:58.030 \longrightarrow 00:25:01.370$ And so, one way to also evaluate
- $516\ 00:25:01.370 --> 00:25:03.180$ these position we call curves is to just look
- 517 00:25:03.180 --> 00:25:05.750 at the area under the curve, which is so A here
- $518\ 00:25:05.750 \longrightarrow 00:25:07.490$ is 0.7 and B's 0.52.
- 519 00:25:07.490 --> 00:25:09.910 And that tells us that on an average
- $520\ 00:25:09.910 \longrightarrow 00:25:14.910$ A ranks edges better as opposed to B.
- $521\ 00:25:16.320 --> 00:25:19.280$ Now, we would like to use near this,
- $522\ 00:25:19.280 \longrightarrow 00:25:22.350$ and the question is what could we use as
- 523 00:25:22.350 --> 00:25:23.183 a gold standard?
- $524\ 00:25:24.030 \longrightarrow 00:25:25.500$ Now, this is real biological data
- 525 00:25:25.500 --> 00:25:28.570 that we are using, and for that,
- 526~00:25:28.570 --> 00:25:31.670 we would also need to look into
- $527~00{:}25{:}31.670 \dashrightarrow 00{:}25{:}35.400$ the literature to find validation.
- $528\ 00:25:35.400 \longrightarrow 00:25:37.460$ So, one good source of information
- $529\ 00:25:37.460 --> 00:25:39.310$ is the escape database curated
- $530\ 00:25:39.310 \longrightarrow 00:25:41.010$ by the Ma'ayan lab.
- $531\ 00:25:41.010 --> 00:25:44.150$ And this database includes the results
- $532\ 00:25:44.150 --> 00:25:46.710$ of loss of function and gain of experiments

- $533\ 00:25:46.710 \longrightarrow 00:25:48.640$ done on genes, and also
- 534 00:25:48.640 --> 00:25:50.010 and also ChIP-seq experiments,
- $535\ 00:25:50.010 \longrightarrow 00:25:51.700$ which identify binding sites
- $536\ 00:25:51.700 --> 00:25:53.193$ of transcription factors.
- 537~00:25:54.330 --> 00:25:57.940 Now, the problem being that even this database
- $538\ 00:25:57.940 --> 00:26:00.970$ is incomplete because the gaps
- $539\ 00:26:00.970 \longrightarrow 00:26:03.980$ in biological knowledge remain and doesn't,
- 540 00:26:03.980 --> 00:26:05.530 I guess over the time, over time,
- 541 00:26:05.530 --> 00:26:08.630 it would be completed, filled more and more.
- 542 00:26:08.630 --> 00:26:12.020 But when we were doing this evaluation,
- 543 00:26:12.020 --> 00:26:14.330 we had to deal with what was effectively
- 544 00:26:14.330 --> 00:26:15.500 a partial gold standard,
- 545 00:26:15.500 --> 00:26:17.760 or an incomplete gold standard.
- $546\ 00:26:17.760 --> 00:26:20.290$ So, the evaluation that we did was not
- 547 00:26:20.290 --> 00:26:22.920 for all of the genes in the dataset,
- $548\ 00:26:22.920 \longrightarrow 00:26:26.453$ but only a fraction of the genes.
- $549\ 00:26:28.210 \longrightarrow 00:26:32.940$ So, we had these two methods
- $550~00{:}26{:}32.940 \dashrightarrow 00{:}26{:}36.470$ and two pseudo times, which we got from that.
- $551\ 00:26:36.470 \longrightarrow 00:26:37.790$ So, what we wanted, what we did
- 552 00:26:37.790 --> 00:26:41.710 is we compared the performance of SINGE
- $553\ 00:26:42.740 \longrightarrow 00:26:46.200$ using say, Monocle 2 and the pseudo time,
- $554\ 00:26:46.200 \longrightarrow 00:26:48.940$ as well as Monocle 2 with only the ordering.
- $555\ 00:26:48.940 --> 00:26:50.290$ And some of the least PAGA Tree
- $556\ 00:26:50.290 \longrightarrow 00:26:51.690$ fed the pseudo time and PAGA Tree
- $557\ 00:26:51.690 \longrightarrow 00:26:52.840$ with only the ordering.
- $558\ 00:26:53.710 \longrightarrow 00:26:57.650$ And so, this is how the precision recall curves
- 559~00:26:57.650 --> 00:27:01.060 of these four methods look.
- $560\ 00:27:01.060 --> 00:27:04.420$ So, we look at the average precision,
- $561\ 00:27:04.420 \longrightarrow 00:27:06.240$ which is the same thing as the area under

- $562\ 00:27:06.240 \longrightarrow 00:27:07.890$ the precision recall curve.
- $563~00{:}27{:}07.890 \dashrightarrow 00{:}27{:}09.783$ And we also look at the average precision
- $564\ 00:27:09.783 \longrightarrow 00:27:13.640$ in the early part of the precision recall curve.
- 565 00:27:13.640 --> 00:27:16.313 And the point for that being that,
- 566 00:27:17.960 --> 00:27:20.650 in say, a usual workflow,
- 567 00:27:20.650 --> 00:27:23.850 you would have a combination method,
- 568 00:27:23.850 --> 00:27:27.713 which would point to some important edges,
- $569\ 00:27:28.620 \longrightarrow 00:27:31.490$ and then, you would potentially tell
- $570\ 00:27:31.490 --> 00:27:33.680$ a collaborator to try
- $571\ 00:27:33.680 \longrightarrow 00:27:35.820$ and experimentally validate that.
- 572 00:27:35.820 --> 00:27:38.190 And in that sense, you would be giving
- $573~00{:}27{:}38.190 \dashrightarrow 00{:}27{:}40.110$ them results from the top of your list,
- 574~00:27:40.110 --> 00:27:43.100 as opposed to trying to tell how well
- $575\ 00:27:43.100 \longrightarrow 00:27:45.300$ the 10,000th edge in the list
- $576\ 00:27:45.300 \longrightarrow 00:27:46.673$ is placed in the rankings.
- $577\ 00:27:47.580 \longrightarrow 00:27:49.820$ So, with that in mind, we also look
- 578 00:27:49.820 --> 00:27:52.780 at what's the average early precision
- $579\ 00:27:52.780 \longrightarrow 00:27:54.093$ of these curves.
- 580 00:27:55.030 --> 00:27:58.540 And for that, we basically say what happened,
- $581\ 00:27:58.540 \longrightarrow 00:28:03.057$ as to what extent is the precision maintained
- $582\ 00:28:04.150 \longrightarrow 00:28:06.380$ until 10% of the genes
- $583\ 00:28:06.380 \longrightarrow 00:28:08.340$ and the gold standard are...
- 584 00:28:08.340 --> 00:28:09.700 Or interactions with the gold standard
- $585\ 00:28:09.700 \longrightarrow 00:28:14.550$ are regarded in the list that we have.
- $586\ 00:28:14.550 \longrightarrow 00:28:17.760$ So, the figure to the right shows
- $587\ 00:28:17.760 \longrightarrow 00:28:19.900$ a scatterplot of these, the average precision
- $588\ 00:28:19.900 \longrightarrow 00:28:21.080$ and the average early precision
- $589\ 00:28:21.080 --> 00:28:24.670$ for these four methods, for these four options.
- $590\ 00:28:24.670 \longrightarrow 00:28:27.420$ And what we see is that the...
- $591\ 00:28:27.420 --> 00:28:29.380$ The best performing combination
- 592 00:28:29.380 --> 00:28:31.170 is using Monocle's ordering,

- 593~00:28:31.170 --> 00:28:35.540 but not its pseudo time, and Monocle applying
- 594 00:28:35.540 --> 00:28:37.610 the pseudo time that it order,
- $595\ 00:28:37.610 \longrightarrow 00:28:41.050$ that it assigns to the cells,
- $596\ 00:28:41.050 \longrightarrow 00:28:44.213$ actually degrades the performance quite a bit.
- 597 00:28:45.670 --> 00:28:48.950 And both of the PAGA Tree options
- 598 00:28:48.950 --> 00:28:50.600 with, or without pseudo time,
- $599\ 00:28:50.600 \longrightarrow 00:28:52.350$ are in between these.
- 600 00:28:52.350 --> 00:28:55.680 So, now why would this happen?
- $601\ 00:28:55.680 \longrightarrow 00:28:57.100$ For example, and let's take
- 602 00:28:57.100 --> 00:28:58.420 an extreme case, right?
- 603 00:28:58.420 --> 00:29:01.630 And okay, before that, there's not necessarily
- 604 00:29:04.060 --> 00:29:05.680 something that's wrong with Monocle,
- 605 00:29:05.680 --> 00:29:08.960 but it's basically that for this dataset,
- $606\ 00:29:08.960 \longrightarrow 00:29:11.530$ in this instance, the pseudo time values
- $607\ 00:29:11.530 \longrightarrow 00:29:14.500$ did not necessarily make a lot of sense.
- 608 00:29:14.500 --> 00:29:17.170 So, let's say you have perfectly ordered cells.
- 609 00:29:17.170 --> 00:29:18.890 And for the first half of the cells,
- 610 00:29:18.890 --> 00:29:22.110 you just assign a value very close
- $611\ 00:29:22.110 \longrightarrow 00:29:23.180$ to zero and the second half,
- $612\ 00:29:23.180 \longrightarrow 00:29:25.570$ you assign a value very close to one.
- 613 00:29:25.570 --> 00:29:27.470 So, even though the ordering of the cells
- 614 00:29:27.470 --> 00:29:31.170 was quite nice and reliable, just because
- $615\ 00:29:31.170 \longrightarrow 00:29:33.810$ we ended up assigning a value
- $616\ 00:29:33.810 \longrightarrow 00:29:35.750$ to the pseudo times, often times,
- $617\ 00:29:35.750 \longrightarrow 00:29:38.090$ which is completely unrealistic.
- $618\ 00:29:38.090 \longrightarrow 00:29:40.970$ We might end up losing
- $619\ 00:29:40.970 \longrightarrow 00:29:41.950$ a lot of information
- $620\ 00:29:41.950 \longrightarrow 00:29:44.320$ that we otherwise had in the dataset,
- $621\ 00:29:44.320 \longrightarrow 00:29:45.270$ or in the ordering.
- $622\ 00:29:48.920 \longrightarrow 00:29:52.880$ So, yeah, as an extended,
- 623 00:29:52.880 --> 00:29:55.520 the ideas from this particular figure, right?

- $624\ 00:29:55.520 \longrightarrow 00:29:57.240$ So, you have two methods,
- 625 00:29:57.240 --> 00:29:59.030 they're giving you two different...
- 626 00:30:00.010 --> 00:30:01.323 Okay, two methods with their orderings
- $627\ 00:30:01.323 --> 00:30:04.760$ and pseudo times, so basically four cases,
- 628 00:30:04.760 --> 00:30:07.590 and they all give you different rankings,
- 629 00:30:07.590 --> 00:30:12.390 which have different performances
- $630\ 00:30:12.390 \longrightarrow 00:30:14.410$ in terms of network evaluation.
- $631\ 00:30:14.410 \longrightarrow 00:30:16.670$ And in a sense, you could say
- $632\ 00{:}30{:}18.711 \dashrightarrow 00{:}30{:}21.960$ that each of these PAGA Tree inference methods
- $633\ 00:30:21.960 \longrightarrow 00:30:24.870$ itself with all their inefficiencies
- 634 00:30:24.870 --> 00:30:27.990 and efficiencies are only partially looking
- $635\ 00:30:27.990 \longrightarrow 00:30:29.990$ at the biological data.
- $636\ 00:30:29.990 \longrightarrow 00:30:33.730$ So, from that perspective, each
- $637\ 00:30:33.730 \longrightarrow 00:30:37.150$ of these orderings and pseudo time values
- $638\ 00:30:37.150 \longrightarrow 00:30:38.560$ can be considered as sources
- 639 00:30:38.560 --> 00:30:39.980 of noisy information,
- $640\ 00:30:39.980 \longrightarrow 00:30:41.580$ or noisy sources of information.
- 641 00:30:42.430 --> 00:30:46.357 So, instead of trying to just infer
- $642\ 00:30:48.860 --> 00:30:52.100$ one pseudo time trajectory from
- 643 00:30:52.100 --> 00:30:54.810 the dataset and finding the network,
- 644 00:30:54.810 --> 00:30:55.940 or say another, and finding
- $645\ 00:30:55.940 \longrightarrow 00:30:58.480$ the network from that, we could think
- $646\ 00:30:58.480 \longrightarrow 00:31:01.380$ of the trajectory inference method itself
- $647\ 00:31:01.380 \longrightarrow 00:31:03.140$ as an additional hyper parameter
- $648\ 00:31:03.140 --> 00:31:06.310$ on top of the sparsity, and kernel bits,
- $649\ 00:31:06.310 \longrightarrow 00:31:07.520$ and so on.
- 650 00:31:07.520 --> 00:31:10.290 So, instead of aggregating at this point
- 651 00:31:10.290 --> 00:31:11.900 after just one trajectory inference method,
- $652\ 00:31:11.900 \longrightarrow 00:31:14.040$ we could just say that maybe
- $653\ 00:31:14.040 \longrightarrow 00:31:16.110$ we have four trajectory inference methods

- $654\ 00:31:18.950 \longrightarrow 00:31:20.200$ in the beginning.
- $655\ 00:31:20.200 \longrightarrow 00:31:22.760$ And after that, we do all
- 656 00:31:22.760 --> 00:31:24.980 of these sub sampling and application
- 657 00:31:24.980 --> 00:31:27.520 of hyper-parameters, and multiple tests.
- $658\ 00:31:27.520 \longrightarrow 00:31:29.370$ And then, we aggregate over all
- $659\ 00:31:29.370 \longrightarrow 00:31:30.910$ of these results across
- $660\ 00:31:30.910 --> 00:31:32.820$ trajectory inference methods.
- 661 00:31:32.820 --> 00:31:34.080 So, hopefully what that would do
- $662\ 00:31:34.080 \longrightarrow 00:31:39.080$ is that would account for all the inefficiencies,
- 663 00:31:39.290 --> 00:31:40.470 or counter then inefficiencies
- 664 00:31:40.470 --> 00:31:42.910 of individual trajectory inference methods,
- $665\ 00{:}31{:}42.910 {\:{\mbox{--}}\!>}\ 00{:}31{:}47.383$ and give us a more robust network at the end.
- $666\ 00:31:49.280 \longrightarrow 00:31:52.110$ And I have not, I guess, shown
- 667 00:31:52.110 --> 00:31:54.010 our comparisons for the other methods,
- 668 00:31:55.730 --> 00:31:57.640 which obviously isn't in our paper.
- $669\ 00:31:57.640 \longrightarrow 00:31:59.760$ We are doing better than them.
- $670\ 00:31:59.760 \longrightarrow 00:32:01.800$ So, but you can have a look at
- 671 00:32:02.680 --> 00:32:05.030 that in the paper if you're interested,
- 672 00:32:05.030 --> 00:32:07.670 because I just wanted to conceptually focus
- $673\ 00:32:07.670 \longrightarrow 00:32:09.723$ on these ideas a little bit more.
- 674 00:32:11.010 --> 00:32:13.900 So, I guess, one problem with trying
- 675 00:32:13.900 --> 00:32:16.570 to run four different, or five different
- $676\ 00:32:16.570 \longrightarrow 00:32:19.060$ trajectory inference methods is depending on
- 677 00:32:19.060 --> 00:32:20.237 what kind of data set you have
- 678 00:32:20.237 --> 00:32:22.393 and what kind of biology you are studying,
- 679 00:32:23.370 --> 00:32:27.330 you might not necessarily have
- $680\ 00:32:27.330 \longrightarrow 00:32:28.850$ to try only four methods.
- $681\ 00:32:28.850 \longrightarrow 00:32:29.683$ You will probably have
- 682 00:32:29.683 --> 00:32:32.080 to try multiple methods before,
- $683\ 00:32:32.080 \longrightarrow 00:32:34.210$ which let's say, if you know
- 684 00:32:34.210 --> 00:32:35.310 it's a branching trajectory,

- 685 00:32:35.310 --> 00:32:38.150 you end up seeing a branching trajectory.
- $686\ 00:32:38.150 \longrightarrow 00:32:41.200$ And each of these methods would have
- 687 00:32:41.200 --> 00:32:44.320 their own input data format,
- 688 00:32:44.320 --> 00:32:46.503 up data formats, visualizations,
- $689\ 00:32:49.267 \longrightarrow 00:32:52.300$ and all of these other intricacies.
- $690\ 00:32:52.300 --> 00:32:55.170$ And that's where the dynverse project comes
- $691\ 00:32:55.170 \longrightarrow 00:32:56.430$ to our rescue.
- $692\ 00:32:56.430 \longrightarrow 00:32:59.790$ So, if anyone is looking to do
- 693 00:32:59.790 --> 00:33:01.000 a lot of trajectory inference methods,
- $694\ 00:33:01.000 \longrightarrow 00:33:03.900$ I would strongly encourage you to look at that.
- $695\ 00:33:03.900 \longrightarrow 00:33:06.000$ So, these in this project,
- 696 00:33:06.000 --> 00:33:09.850 they have streamlined the use of, I think,
- $697\ 00:33:09.850 \longrightarrow 00:33:11.670\ 55\ trajectory\ inference\ methods.$
- 698 00:33:11.670 --> 00:33:14.060 So, you don't necessarily need to install
- $699\ 00:33:14.060 \longrightarrow 00:33:14.893$ each one of them.
- 700 00:33:14.893 --> 00:33:16.300 You just install this project
- $701\ 00:33:16.300 \longrightarrow 00:33:18.240$ and they run each
- $702\ 00:33:18.240 \longrightarrow 00:33:20.700$ of these methods using a docker.
- $703\ 00:33:20.700 --> 00:33:23.470$ And so, what it also helps you do
- 704 00:33:23.470 --> 00:33:26.160 is it helps you visualize
- $705\ 00:33:26.160 \longrightarrow 00:33:31.160$ all of these trajectories and evaluate them using
- $706\ 00:33:31.420 --> 00:33:34.590$ the same, I guess, support scripts
- 707 00:33:34.590 --> 00:33:37.900 and support functions, which they also provide
- $708\ 00:33:37.900 \longrightarrow 00:33:41.720$ And in all this, this would make
- $709\ 00:33:41.720 \longrightarrow 00:33:43.920$ your lives quite easy.
- $710\ 00:33:43.920 \longrightarrow 00:33:46.850$ And they also have basically a user,
- 711 00:33:46.850 --> 00:33:48.020 a graphical user interface,
- 712 00:33:48.020 --> 00:33:51.740 which helps you prioritize
- 713 00:33:51.740 --> 00:33:55.060 what trajectory inference method to use,

- 714 00:33:55.060 --> 00:33:59.902 depending on what biology you want to study.
- 715 00:33:59.902 --> 00:34:02.340 How many cells you have, what compute power
- $716\ 00:34:02.340 \longrightarrow 00:34:05.973$ you might have access to, and so on.
- 717 00:34:12.272 --> 00:34:16.650 So, okay just some final comments on the use
- 718 00:34:16.650 --> 00:34:18.980 of, I guess, the utility of trajectory inference
- $719\ 00:34:18.980 \longrightarrow 00:34:22.250$ and pseudo times for further analysis.
- 720 00:34:22.250 --> 00:34:24.980 And so, first of all, as in trajectories
- 721 00:34:24.980 --> 00:34:27.990 look really nice, they visually,
- $722\ 00:34:27.990 \longrightarrow 00:34:30.510$ they give us a lot of information.
- 723 00:34:30.510 --> 00:34:33.270 And so, based on what we saw,
- $724\ 00:34:33.270 \longrightarrow 00:34:35.773$ we did see that there's some,
- $725\ 00:34:38.646 \longrightarrow 00:34:40.510$ the ordering information
- $726\ 00:34:40.510 --> 00:34:43.040$ and the pseudo time values can help
- $727\ 00:34:43.040 \longrightarrow 00:34:44.113$ in network inference.
- 728 00:34:45.090 --> 00:34:48.520 The good pseudo times can help a little bit,
- $729\ 00:34:48.520 \longrightarrow 00:34:51.480$ but if you have exceptionally bad pseudo times,
- 730 00:34:51.480 --> 00:34:54.033 it can hurt a lot as opposed to ordering.
- 731 00:34:54.960 --> 00:34:58.710 And not every dataset is really suitable
- 732 00:34:58.710 --> 00:34:59.560 for trajectory inference.
- 733 $00:34:59.560 \longrightarrow 00:35:00.820$ What do I mean by that?
- 734 00:35:00.820 --> 00:35:04.160 So, the dataset that I chose,
- $735\ 00:35:04.160 \longrightarrow 00:35:07.430$ and I guess a lot of what is...
- $736\ 00:35:08.400 --> 00:35:09.630$ What particular inference methods
- $737\ 00:35:09.630 \longrightarrow 00:35:11.190$ are built around, as say,
- 738 00:35:11.190 --> 00:35:14.310 stem cell differentiation in general,
- $739\ 00:35:14.310 \longrightarrow 00:35:19.220$ where it's as in the biology is quite neat
- $740\ 00:35:19.220 \longrightarrow 00:35:20.053$ to begin with.
- 741 00:35:20.053 --> 00:35:23.260 As in you start off from a single cell type,
- $742\ 00:35:23.260 \longrightarrow 00:35:27.180$ and a lot of the biology is already known.

```
743 00:35:27.180 --> 00:35:30.010 So, you don't have to worry, you know
```

- 744 00:35:30.010 --> 00:35:31.820 that it's going to be a branching,
- 745 00:35:31.820 --> 00:35:36.460 or bifurcating, or multi furcating trajectory.
- 746 00:35:36.460 --> 00:35:38.130 So, you know that the quality of the biology,
- 747 00:35:38.130 --> 00:35:42.580 you know what cell states to exist, to expect,
- $748\ 00:35:42.580 \longrightarrow 00:35:43.770$ and so on, and so forth.
- $749\ 00:35:43.770 --> 00:35:45.930$ You know the markers of each of those.
- 750 00:35:45.930 --> 00:35:49.360 And so, studying something like that
- $751\ 00:35:49.360 \longrightarrow 00:35:53.040$ is much more easier using trajectory inference,
- 752 00:35:53.040 --> 00:35:54.460 or pseudo time.
- $753\ 00:35:54.460 \longrightarrow 00:35:55.970$ On the other hand, let's say,
- $754\ 00:35:55.970 --> 00:35:59.330$ if you had a sample from a cancer tumor
- $755\ 00:35:59.330 \longrightarrow 00:36:02.380$ in that you would find cancer cells,
- 756 00:36:02.380 --> 00:36:06.140 normal cells, a bunch of immune cells,
- 757 00:36:06.140 --> 00:36:10.350 probably 10 to 20 kinds of immune cells,
- $758\ 00:36:10.350 \longrightarrow 00:36:11.580$ and so on.
- 759 00:36:11.580 --> 00:36:13.620 So, the trajectory inference method
- 760 00:36:14.680 --> 00:36:18.040 usually tracks, or predicts places,
- $761\ 00:36:18.040 \longrightarrow 00:36:19.820$ cell states and context.
- $762\ 00:36:19.820 \longrightarrow 00:36:22.770$ Not cell types themselves.
- $763\ 00:36:22.770 \longrightarrow 00:36:25.270$ So, you wouldn't necessarily be able
- $764\ 00:36:25.270 -> 00:36:28.560$ to reliably run a trajectory inference method
- 765 00:36:28.560 --> 00:36:33.020 across as in using a mix of different cell types,
- $766\ 00:36:33.020 \longrightarrow 00:36:34.990$ as opposed to cell states.
- 767 00:36:34.990 --> 00:36:38.220 Now, with the stem cell differentiation,
- $768\ 00:36:38.220 \longrightarrow 00:36:40.780$ the good thing is that the cell states
- $769\ 00:36:40.780 \longrightarrow 00:36:43.460$ themselves after a point, transition
- 770 00:36:43.460 --> 00:36:45.200 into different cell types,
- 771 $00:36:45.200 \longrightarrow 00:36:47.030$ because it's the same cell,
- 772 00:36:47.030 --> 00:36:50.170 or same cell type which transitions
- 773 00:36:50.170 --> 00:36:51.803 through multiple cell types,

- $774\ 00:36:52.800 \longrightarrow 00:36:55.560$ through these cell states.
- 775 00:36:55.560 --> 00:36:58.170 But that's not the case with cancer biology,
- $776\ 00:36:58.170 \longrightarrow 00:37:00.640$ where you already start off
- 777 00:37:00.640 --> 00:37:05.640 with a mix of cell types and trajectory inference
- $778\ 00:37:05.800 \longrightarrow 00:37:08.430$ would not make sense for that mix.
- 779 00:37:08.430 --> 00:37:10.610 What people have tried is isolate,
- 780 00:37:10.610 --> 00:37:15.610 just say a T-cell type, and then try
- $781\ 00:37:15.910 \longrightarrow 00:37:18.920$ to order, or find the trajectory only
- $782\ 00:37:18.920 \longrightarrow 00:37:21.230$ for those T-cells.
- $783\ 00:37:21.230 \longrightarrow 00:37:23.350$ And there has been some success in that.
- $784\ 00:37:23.350 \longrightarrow 00:37:27.230$ So, you could run trajectory inference
- $785\ 00:37:27.230 --> 00:37:29.920$ for a subset of the dataset, but not necessarily
- $786\ 00:37:29.920 \longrightarrow 00:37:30.870$ the entire dataset.
- 787 00:37:32.230 --> 00:37:37.230 And so, depending on what biological processes
- $788\ 00:37:37.910 \longrightarrow 00:37:38.853$ you want to study,
- $789\ 00:37:40.820 \longrightarrow 00:37:42.620$ there are trajectory inference methods,
- 790 00:37:42.620 --> 00:37:44.600 which may or may not be suitable for it.
- 791 00:37:44.600 --> 00:37:47.350 For example, a number of methods
- 792 00:37:47.350 --> 00:37:51.140 like Monocle and PAGA Tree,
- 793 00:37:51.140 --> 00:37:56.140 they try to find tree-like structures
- $794\ 00:37:56.300 \longrightarrow 00:37:58.180$ in the trajectories,
- $795\ 00:37:58.180 \longrightarrow 00:37:59.230$ so they would not be suitable
- 796 00:37:59.230 --> 00:38:03.440 for a cyclic biological process
- $797\ 00:38:03.440 \longrightarrow 00:38:06.043$ like just maintenance processes in cells.
- $798\ 00:38:06.900 \longrightarrow 00:38:08.010$ And then, there are other methods
- 799 00:38:08.010 --> 00:38:10.800 which actually try to find cell cycles,
- $800\ 00:38:10.800 \longrightarrow 00:38:12.210$ and they would not be appropriate
- $801\ 00:38:12.210 --> 00:38:13.653$ for branching processes.
- 802 00:38:15.550 --> 00:38:18.630 And I guess, as a no single

- 803 00:38:18.630 --> 00:38:20.030 trajectory inference method,
- 804 00:38:22.550 --> 00:38:25.200 accurately represents the biology.
- 805 00:38:25.200 --> 00:38:26.720 So, it's all basically
- $806\ 00:38:26.720 --> 00:38:28.780$ some mathematical abstraction
- $807\ 00:38:28.780 \longrightarrow 00:38:31.133$ of what might be happening in the cells.
- $808\ 00:38:35.216 \longrightarrow 00:38:36.049$ And yeah, as an if...
- $809\ 00:38:36.049 \longrightarrow 00:38:37.170$ If at the outset, you know
- 810 00:38:37.170 --> 00:38:41.090 what kind of trajectory to expect, then it helps
- 811 00:38:41.090 --> 00:38:42.240 in trying to
- $812\ 00:38:44.770 \longrightarrow 00:38:45.910$ at least first really,
- 813 00:38:45.910 --> 00:38:49.790 say whether the trajectory that you're getting
- 814 00:38:49.790 --> 00:38:52.060 and the pseudo times that you get
- $815\ 00:38:52.060 \longrightarrow 00:38:55.030$ is of any worth.
- 816 00:38:55.030 --> 00:38:57.920 So, just to give you an example.
- $817\ 00:38:57.920 \longrightarrow 00:39:00.290$ So, we started off with Monocle 2
- $818\ 00:39:00.290 \longrightarrow 00:39:02.560$ as one of our examples in our paper,
- $819\ 00:39:02.560 \longrightarrow 00:39:05.330$ and then we wanted to have another method
- $820\ 00:39:05.330 \longrightarrow 00:39:07.413$ to compare the effects of different
- 821 00:39:07.413 --> 00:39:09.890 trajectory inference methods.
- $822\ 00:39:09.890 --> 00:39:12.850$ And PAGA Tree was not necessarily the first one.
- $823\ 00:39:12.850 \longrightarrow 00:39:14.290$ We tried a number of other ones,
- $824\ 00:39:14.290 \longrightarrow 00:39:16.080$ which did not.
- $825\ 00:39:16.080 \longrightarrow 00:39:18.380$ And we knew what to expect here.
- 826 00:39:18.380 --> 00:39:21.160 We knew that there was stem cell
- 827 00:39:21.160 --> 00:39:26.160 to ectoderm trajectory and endoderm trajectory,
- $828\ 00:39:26.430 \longrightarrow 00:39:27.980$ or a branch of that.
- 829 00:39:27.980 --> 00:39:32.980 And using basically, just the first,
- $830\ 00:39:35.040 \longrightarrow 00:39:37.560$ I think we tried four methods

- 831 $00:39:37.560 \longrightarrow 00:39:39.400$ and PAGA Tree was basically the fourth method,
- $832\ 00:39:39.400 \longrightarrow 00:39:41.690$ which gave us that kind of branching trajectory,
- 833 00:39:41.690 --> 00:39:44.870 or branching topology for the biology.
- 834 00:39:44.870 --> 00:39:49.250 And so, none of the methods you try
- 835 00:39:49.250 --> 00:39:51.803 might necessarily mean anything,
- $836\ 00:39:53.010 \longrightarrow 00:39:55.503$ unless you have some way of validating that.
- 837 00:39:56.520 --> 00:39:59.010 So, at this point, I'm gonna switch
- 838 00:39:59.010 --> 00:40:04.010 to spatial expression,
- $839\ 00:40:04.070 \longrightarrow 00:40:05.820$ or a spatial data and special analysis.
- $840\ 00:40:05.820 \longrightarrow 00:40:08.230$ So, if you have any questions
- 841 00:40:08.230 --> 00:40:12.380 about the pseudo time analysis,
- $842\ 00:40:12.380 \longrightarrow 00:40:13.813$ should we take it now, or?
- 843 00:40:19.010 --> 00:40:20.270 <v Lecturer>Does any body have any questions</v>
- 844 00:40:20.270 --> 00:40:22.753 on the first half of the presentation here?
- 845 00:40:26.265 --> 00:40:27.290 <v Dr. Deshpande>Oh, we can continue on,</v>
- $846\ 00:40:27.290 \longrightarrow 00:40:29.707$ then we can come back later.
- $847\ 00:40:34.439 \longrightarrow 00:40:35.689$ Shall we go on?
- 848 00:40:40.670 --> 00:40:42.148 <v Lecturer>Sounds good.</v>
- $849\ 00:40:42.148 \longrightarrow 00:40:44.148 < v Dr. Deshpande>Okay.</v>$
- $850\ 00:40:47.730 \longrightarrow 00:40:49.653$ Okay, so that was all about,
- $851\ 00:40:51.940 \longrightarrow 00:40:56.940$ say how pseudo time is used in our analysis.
- 852 00:40:56.940 --> 00:41:01.617 And so, the other end of,
- $853\ 00:41:03.170 \longrightarrow 00:41:04.470\ I$ guess, not necessarily end,
- $854\ 00:41:04.470 \longrightarrow 00:41:05.310$ the other perspective
- 855 00:41:05.310 --> 00:41:09.600 is how is space important and how,
- 856 00:41:09.600 --> 00:41:11.050 what kind of data do we have,
- 857 00:41:12.540 --> 00:41:14.843 which give us information about space?
- $858\ 00:41:15.760 \longrightarrow 00:41:18.450$ So, the spatial context of cells

```
859\ 00:41:18.450 --> 00:41:21.570 is very important in many biological processes.
```

- $860~00{:}41{:}21.570 \dashrightarrow 00{:}41{:}23.690$ For example, when immune cells respond
- 861 00:41:23.690 --> 00:41:26.670 to an infection, or a wound, they need
- 862 00:41:26.670 --> 00:41:28.920 to be in physical proximity of their targets.
- 863 00:41:31.010 --> 00:41:33.840 Similarly with, I guess, cancer tumor growth,
- $864\ 00:41:33.840 \longrightarrow 00:41:37.560$ and the immune response to cancer
- $865\ 00:41:37.560 \longrightarrow 00:41:39.720$ happen through intracellular signaling.
- 866 00:41:39.720 --> 00:41:42.390 Either through cytokine secretion,
- $867\ 00:41:42.390 \longrightarrow 00:41:45.563$ or through surface receptors on adjacent cells.
- 868 00:41:48.410 --> 00:41:50.170 Just knowing the relative location
- $869\ 00:41:50.170 \longrightarrow 00:41:51.520$ of different cell types can also
- $870\ 00:41:51.520 \longrightarrow 00:41:52.820$ be very informative.
- 871 00:41:52.820 --> 00:41:55.393 For example, in the figure here,
- 872 00:41:56.630 --> 00:41:57.960 the information about the presence
- 873 00:41:57.960 --> 00:42:01.920 of various immune cell types nearest tumor,
- $874\ 00:42:01.920 \longrightarrow 00:42:03.720$ and the extent of immune deficient
- $875\ 00:42:03.720 \longrightarrow 00:42:06.163$ in the tumor are essential prognostic markers.
- 876 00:42:07.930 --> 00:42:10.850 And so, single cell RNA-seq,
- 877 00:42:12.550 --> 00:42:15.270 as good as it is, it associates a cell
- $878\ 00:42:15.270 \longrightarrow 00:42:17.900$ from its tissue, due to which
- $879\ 00:42:17.900 \longrightarrow 00:42:20.820$ we lose the spatial context of the cell states.
- $880\ 00:42:20.820 \longrightarrow 00:42:23.580$ But in recent years, we have been able
- $881\ 00:42:23.580 \longrightarrow 00:42:28.170$ to develop both
- 882 00:42:28.170 --> 00:42:29.760 as in spatial proteomics,
- $883\ 00:42:29.760 \longrightarrow 00:42:34.760$ which help you to image protein
- $884\ 00:42:35.080 \longrightarrow 00:42:39.800$ and densities of say, up to 30 markers
- $885\ 00:42:39.800 \longrightarrow 00:42:41.883$ at single cell resolution in the tissue.
- 886 00:42:43.340 --> 00:42:45.910 As well as spatial transcriptomics,
- $887\ 00:42:45.910 \longrightarrow 00:42:50.540$ which can measure 20,000 genes at spots
- $888\ 00:42:50.540 \longrightarrow 00:42:52.970$ in the tissue.

```
889 00:42:52.970 --> 00:42:56.610 And this was named method of the year last year
```

- $890\ 00:42:56.610 \longrightarrow 00:42:58.883$ in 2020, yeah, that was last year.
- $891\ 00:43:01.071 \longrightarrow 00:43:04.580$ So, here's just a workflow
- 892 00:43:04.580 --> 00:43:06.450 of the next Visium technology,
- $893\ 00:43:06.450 \longrightarrow 00:43:07.283$ which is one of these
- $894\ 00:43:07.283 \longrightarrow 00:43:09.950$ spatial transcriptomics technologies.
- $895\ 00:43:09.950 \longrightarrow 00:43:14.853$ So, this includes 5,000 barcoded spots on slide.
- $896\ 00:43:15.800 \longrightarrow 00:43:20.800$ And these are added to the cells in the...
- $897\ 00:43:21.250 \longrightarrow 00:43:24.110$ Which are located in those spots.
- 898 00:43:24.110 --> 00:43:26.120 And this helps preserve the spatial context
- $899\ 00:43:26.120 \longrightarrow 00:43:28.533$ of the cells to the actual sequencing.
- 900 00:43:29.600 --> 00:43:33.240 Now, this technology is not exactly single cell.
- 901 00:43:33.240 --> 00:43:37.363 It still provides a lot of useful spacial detail.
- 902 00:43:41.080 --> 00:43:46.080 So yeah, for explaining this project,
- 903 00:43:46.860 --> 00:43:50.770 I will use the 10x Visium sample,
- $904\ 00:43:50.770 \longrightarrow 00:43:52.140$ provided by 10x genomics
- $905\ 00:43:53.040 \longrightarrow 00:43:54.950$ of a breast cancer tissue.
- 906 00:43:54.950 --> 00:43:56.700 So, the figure on the left
- 907 00:43:56.700 --> 00:43:59.690 is an H and E slide, it's hematoxylin
- $908\ 00:44:00.785 \longrightarrow 00:44:02.580$ and eosin stain slide,
- 909 00:44:02.580 --> 00:44:07.580 which helps pathologists annotate
- 910 $00:44:07.700 \longrightarrow 00:44:12.700$ the sample for tumor, and lesions, and so on.
- 911 00:44:13.680 --> 00:44:18.500 And the second image is that slide annotated
- $912\ 00:44:18.500 \longrightarrow 00:44:21.500$ by a pathologist, and you can see
- 913 00:44:21.500 --> 00:44:24.640 that there are different biology's
- 914 00:44:24.640 --> 00:44:27.130 in this one slide.
- 915 00:44:27.130 --> 00:44:29.090 And for example, the lesion on top
- $916\ 00:44:29.090 \longrightarrow 00:44:31.050$ is an invasive cancer lesion, which means
- 917 00:44:31.050 --> 00:44:33.330 that it can spread beyond the breast tissue,
- 918 00:44:33.330 --> 00:44:34.900 but the other lesions correspond

- 919 $00:44:34.900 \longrightarrow 00:44:35.970$ to DCAs lesions,
- 920 00:44:35.970 --> 00:44:38.840 which are not yet classified as invasive,
- $921\ 00:44:38.840 \longrightarrow 00:44:41.700$ they could in the future be invasive.
- $922\ 00:44:41.700 --> 00:44:43.390$ Other important annotations are those
- $923\ 00:44:43.390 \longrightarrow 00:44:46.600$ of immune cells and the stromal cells
- $924\ 00:44:46.600 \longrightarrow 00:44:48.013$ in between these lesions.
- 925 00:44:49.680 --> 00:44:51.800 For a good clinical outcome, you would hope
- $926\ 00:44:51.800 \longrightarrow 00:44:54.683$ that immune cells can infiltrate these lesions.
- $927\ 00:44:55.590 \longrightarrow 00:44:59.100$ And so the figure on the right shows
- 928 00:44:59.100 \rightarrow 00:45:00.360 the same H and E slide
- $929\ 00:45:02.000 \longrightarrow 00:45:04.660$ with overlaid Visium spots.
- 930 00:45:04.660 --> 00:45:06.870 So, each of these spots correspond
- 931 $00:45:06.870 \longrightarrow 00:45:08.133$ to one measurement.
- $932\ 00:45:09.510 \longrightarrow 00:45:14.400$ So, this slide shows a couple of examples
- 933 $00:45:14.400 \longrightarrow 00:45:16.900$ of spacial gene expression.
- $934\ 00:45:16.900 \longrightarrow 00:45:18.550$ So, the figure to the left
- 935 00:45:18.550 --> 00:45:21.230 is the same annotated H and E slide
- 936 $00:45:21.230 \longrightarrow 00:45:23.480$ that will help us keep track
- $937\ 00:45:23.480 \longrightarrow 00:45:26.870$ of the biology in the slide.
- 938 00:45:26.870 --> 00:45:29.900 And so, the first figure, the middle figure,
- 939 00:45:29.900 --> 00:45:32.770 basically it shows the expression of CD8A,
- 940 00:45:32.770 --> 00:45:35.600 which is a marker of cytotoxic T-cells.
- 941 00:45:35.600 --> 00:45:37.020 Now, we see this gene expressed
- $942\ 00{:}45{:}37.020 \dashrightarrow 00{:}45{:}42.020$ in the blood near the invasive and DCAs lesions,
- $943\ 00:45:42.480 \longrightarrow 00:45:43.810$ which means that the immune cells
- $944\ 00:45:43.810 \longrightarrow 00:45:44.900$ are responding to a tumor.
- $945\ 00:45:44.900 \longrightarrow 00:45:46.550$ However, we see that
- $946\ 00:45:46.550 \longrightarrow 00:45:48.740$ there's not much infiltration of these cells
- $947\ 00:45:48.740 \longrightarrow 00:45:49.743$ within the lesions.
- $948\ 00:45:50.920 \longrightarrow 00:45:53.990$ The second marker is CD14, which is found

```
949 00:45:53.990 --> 00:45:55.740 in macrophages and dendritic cells,
```

- 950 00:45:56.630 --> 00:45:58.310 and its expression is much higher
- 951 00:45:58.310 --> 00:45:59.730 inside the lesions, which could point
- $952\ 00:45:59.730 \longrightarrow 00:46:03.670$ to successful infiltration of these cell types.
- $953\ 00:46:03.670 \longrightarrow 00:46:08.240\ Now$, just a reminder, these the measurements
- $954\ 00:46:08.240 \longrightarrow 00:46:09.870$ that we get from 10x Visium
- 955 00:46:09.870 --> 00:46:13.000 are not exactly single cell, but they're near,
- 956 00:46:13.000 --> 00:46:14.660 near single cell.
- 957 00:46:14.660 --> 00:46:16.490 In a sense that each of these spots
- $958\ 00:46:16.490 \longrightarrow 00:46:18.433$ is 55 micro meters wide.
- 959 00:46:19.360 --> 00:46:21.690 And depending on what cell type
- 960 00:46:21.690 --> 00:46:22.910 you might have in that spot,
- $961\ 00:46:22.910 \longrightarrow 00:46:26.920$ it could have anywhere from one to 10 cells.
- 962 00:46:26.920 --> 00:46:28.140 And immune cells are much smaller,
- 963 00:46:28.140 --> 00:46:30.300 so there could be up to 10 immune cells in it,
- 964 00:46:30.300 --> 00:46:32.280 but maybe only one cancer,
- $965\ 00:46:32.280 \longrightarrow 00:46:34.750$ or epithelial cell in that spot.
- 966 00:46:34.750 --> 00:46:36.410 So, as a result of gene expression
- $967\ 00:46:36.410 \longrightarrow 00:46:38.460$ of that spot is the average
- $968\ 00:46:38.460 \longrightarrow 00:46:39.610$ of the cells inside it.
- 969 00:46:41.910 --> 00:46:46.754 Now, our lab has a method called CoGAPS,
- 970 00:46:46.754 --> 00:46:49.640 oesophageal CoGAPS, which is a Bayesian
- 971 00:46:49.640 --> 00:46:51.480 Markov chain Monte Carlo method
- 972 $00:46:51.480 \longrightarrow 00:46:53.330$ for nonnegative matrix factorization.
- 973 00:46:54.370 --> 00:46:58.470 And so, as a result of say,
- 974 00:46:58.470 --> 00:47:00.940 the 10x Visium measurement,
- $975\ 00:47:00.940 \longrightarrow 00:47:04.210$ we now have a high dimensional matrix
- $976\ 00:47:04.210 \longrightarrow 00:47:08.020$ with 20,000 genes and around 5,000 spots.
- $977\ 00:47:08.020 \longrightarrow 00:47:11.610$ And what CoGAPS does is it helps
- 978 00:47:11.610 --> 00:47:14.730 to factorize this matrix
- 979 00:47:14.730 --> 00:47:17.750 into two low rank matrices,

- 980 00:47:17.750 --> 00:47:19.363 both of which are non-negative,
- 981 00:47:20.860 --> 00:47:25.450 which correspond to latent patterns in the data.
- $982\ 00:47:25.450 \longrightarrow 00:47:26.790$ And in the past, we have seen
- 983 00:47:26.790 --> 00:47:30.150 that these two correspond to biology's
- $984\ 00:47:31.240 \longrightarrow 00:47:33.033$ based on the pattern markers.
- 985 00:47:34.220 --> 00:47:37.780 So, the two matrices that CoGAPS factorizes
- 986 00:47:37.780 --> 00:47:40.500 the dataset into are the amplitude matrix,
- $987\ 00:47:40.500 \longrightarrow 00:47:44.220$ which has say, 20,000 rows for 20,000 genes
- 988 00:47:44.220 --> 00:47:46.573 and N columns for the end patterns.
- 989 00:47:47.560 --> 00:47:49.710 And this helps us identify groups
- 990 00:47:49.710 --> 00:47:51.640 of co-expressed genes,
- 991 00:47:51.640 --> 00:47:53.580 which correspond to the patterns.
- 992 00:47:53.580 --> 00:47:56.520 And the pattern matrix has N rows
- $993~00{:}47{:}56.520 \dashrightarrow 00{:}48{:}01.180$ and $5{,}000$ columns, and they associate the spots
- 994 00:48:01.180 \rightarrow 00:48:03.820 on the sample with patterns.
- 995 00:48:03.820 --> 00:48:07.744 So, because of the nature of the CoGAPS,
- 996 00:48:07.744 --> 00:48:10.850 factorization, and these, the columns
- 997 00:48:10.850 --> 00:48:13.810 of the matrices here, or the rows of the matrices
- 998 00:48:13.810 --> 00:48:15.390 here are not really orthogonal.
- 999 00:48:15.390 --> 00:48:17.410 They are independent, but not orthogonal.
- 1000 00:48:17.410 --> 00:48:20.240 So, they could co-exist in spots,
- $1001\ 00{:}48{:}20.240 \dashrightarrow 00{:}48{:}24.590$ or a gene could be present in multiple processes,
- $1002\ 00:48:24.590 \longrightarrow 00:48:25.423$ and multiple patterns,
- 1003 00:48:25.423 --> 00:48:27.133 which correspond to processes.
- $1004\ 00:48:28.720 \longrightarrow 00:48:33.720$ So, when we apply CoGAPS to the Visium data,
- $1005\ 00:48:36.950 \longrightarrow 00:48:39.050$ so the first try was basically
- $1006\ 00:48:39.050 \longrightarrow 00:48:42.620$ just five patterns, and when we apply it

- $1007\ 00:48:42.620 \longrightarrow 00:48:45.000$ to try and find five patterns
- 1008 00:48:47.390 --> 00:48:50.060 after a factorization, we see that
- $1009\ 00:48:50.060 \longrightarrow 00:48:51.420$ a number of them correspond
- $1010\ 00:48:51.420 \longrightarrow 00:48:53.990$ to the pathology annotations
- $1011\ 00:48:53.990 \longrightarrow 00:48:56.860$ that we see on the figure on the left.
- 1012 00:48:56.860 --> 00:48:58.870 So, we find a pattern which corresponds
- $1013\ 00:48:58.870 \longrightarrow 00:49:01.220$ to the immune cells.
- $1014\ 00:49:01.220 \longrightarrow 00:49:03.710$ We find a pattern which corresponds
- $1015\ 00:49:03.710 --> 00:49:06.900$ to invasive carcinoma on the top left here.
- $1016\ 00:49:06.900 \longrightarrow 00:49:08.640$ And we also find a pattern which corresponds
- $1017\ 00:49:08.640 \longrightarrow 00:49:10.193$ to the DCAs lesions.
- $1018\ 00:49:12.260 \longrightarrow 00:49:14.860$ And as we increase the dimensionality
- 101900:49:14.860 --> 00:49:17.670 of CoGAPS factorization, we start seeing more
- $1020\ 00:49:17.670 \longrightarrow 00:49:19.520$ and more tissue heterogeneity.
- $1021\ 00:49:19.520 \longrightarrow 00:49:23.320$ For example, we now see three patterns
- $1022\ 00{:}49{:}23.320 \dashrightarrow 00{:}49{:}25.580$ which are associated with the mesial carcinoma,
- $1023\ 00:49:25.580 \longrightarrow 00:49:27.160$ and we can see that they correspond
- $1024\ 00:49:27.160 \longrightarrow 00:49:31.440$ to different regions in that lesion.
- $1025\ 00:49:31.440 \longrightarrow 00:49:33.310$ And this for example is completely internal,
- 1026 00:49:33.310 --> 00:49:35.823 which has no interaction with immune cells.
- 1027 00:49:36.680 --> 00:49:38.830 We have a pattern which corresponds
- $1028\ 00:49:38.830 \longrightarrow 00:49:40.620$ to immune cells, we have a pattern
- $1029\ 00:49:40.620 \longrightarrow 00:49:43.350$ which corresponds to the stromal cells.
- $1030\ 00:49:43.350 --> 00:49:46.870$ And we also have different patterns
- $1031\ 00:49:46.870 \longrightarrow 00:49:50.640$ which highlight individual DCAs lesions.
- 1032 00:49:50.640 --> 00:49:53.300 So, one could say that potentially it's trying,
- 1033 00:49:53.300 --> 00:49:56.880 it is finding biology's,
- $1034\ 00:49:56.880 \longrightarrow 00:49:59.273$ which are unique to these DCAs lesions.
- $1035\ 00:50:03.938 \longrightarrow 00:50:06.780$ So, we can analyze the A matrix

```
1036~00{:}50{:}06.780 \dashrightarrow 00{:}50{:}08.610 to identify groups of genes associated
```

- $1037\ 00:50:08.610 \longrightarrow 00:50:10.590$ with each pattern, and we call these
- $1038\ 00:50:10.590 \longrightarrow 00:50:12.030$ the pattern markers.
- 1039 00:50:12.030 --> 00:50:14.590 And these help us identify pathways
- 1040 00:50:14.590 --> 00:50:17.819 that are likely expressed in these patterns,
- 1041 00:50:17.819 --> 00:50:20.350 or because now, especially in this sample,
- $1042\ 00:50:20.350 \longrightarrow 00:50:22.640$ we see a one to one association
- $1043\ 00:50:22.640 --> 00:50:25.290$ between the pattern and the biology,
- $1044\ 00:50:25.290 \longrightarrow 00:50:27.003$ also in the biology, basically.
- $1045\ 00:50:28.630 \longrightarrow 00:50:31.733$ So, let's see, how long do we have.
- $1046\ 00:50:34.649 \longrightarrow 00:50:37.570$ I think we're close to...
- 1047 00:50:37.570 --> 00:50:39.850 I'll quickly rush through these.
- $1048\ 00:50:39.850 \longrightarrow 00:50:44.850$ So, the other analysis that we can do is given,
- $1049\ 00:50:45.480 \longrightarrow 00:50:48.910$ let's say two of these patterns,
- $1050\ 00:50:48.910 --> 00:50:52.390$ we can try to see how these patterns interact.
- $1051\ 00:50:52.390 \longrightarrow 00:50:53.850$ So, you can see that these patterns
- $1052\ 00:50:53.850 \longrightarrow 00:50:57.610$ have a lot of spatial structure to it,
- 1053 00:50:57.610 --> 00:50:59.100 which CoGAPS was not told about.
- $1054~00{:}50{:}59.100$ --> $00{:}51{:}00.840$ CoGAPS, the parameters that Co-GAPS uses
- 1055 00:51:00.840 --> 00:51:02.790 have no special information,
- $1056\ 00:51:02.790 \longrightarrow 00:51:05.730$ and it's still found these spatial structures.
- $1057\ 00:51:05.730 --> 00:51:08.150$ So, and we also see that these patterns
- $1058\ 00:51:08.150 \longrightarrow 00:51:09.510$ are adjacent to each other and we want
- $1059\ 00:51:09.510 \longrightarrow 00:51:11.450$ to see how they interact.
- $1060\ 00:51:11.450 \longrightarrow 00:51:13.033$ So, what we do is we find,
- $1061\ 00:51:14.880 \longrightarrow 00:51:18.430$ basically we estimate the kernel density
- $1062\ 00:51:18.430 --> 00:51:21.910$ of each of these patterns, which is a function
- $1063\ 00:51:21.910 \longrightarrow 00:51:24.630$ of both the pattern intensity at a spot,
- $1064\ 00:51:24.630 \longrightarrow 00:51:28.040$ as well as the spatial clustering
- $1065\ 00:51:28.040 \longrightarrow 00:51:30.400$ of hyper intensities.

```
1066\ 00:51:30.400 \longrightarrow 00:51:31.740 And we compare that against
```

- 1067 00:51:31.740 --> 00:51:34.820 another distribution obtained by
- 1068 00:51:34.820 --> 00:51:37.120 the density estimation after randomizing
- $1069\ 00:51:37.120 --> 00:51:39.353$ the locations of these pattern densities.
- $1070\ 00:51:40.330 \longrightarrow 00:51:43.490$ So, the intensities which are beyond
- 1071 00:51:43.490 --> 00:51:46.570 distal distribution are the ones that we...
- $1072\ 00:51:48.100 \longrightarrow 00:51:49.090$ Are the spots which correspond
- $1073\ 00:51:49.090 \longrightarrow 00:51:50.930$ to these outliers are the ones
- $1074\ 00:51:50.930 \longrightarrow 00:51:55.400$ that we count as hotspots of pattern activity.
- $1075\ 00:51:55.400 \longrightarrow 00:51:57.130$ Similarly, we can find the hotspots
- $1076\ 00:51:57.130 \longrightarrow 00:51:59.380$ of immune response.
- 1077 00:51:59.380 --> 00:52:02.110 And when we combine both of them,
- 1078 00:52:02.110 --> 00:52:07.110 we find regions where cancer is active,
- 1079 00:52:08.720 --> 00:52:10.810 regions where immune cells are active,
- $1080\ 00:52:10.810 \longrightarrow 00:52:13.970$ and regions where both of them are active.
- $1081\ 00:52:13.970 \longrightarrow 00:52:16.090$ And this is the interaction region.
- $1082\ 00:52:16.090 \longrightarrow 00:52:17.490$ And in this region, we are trying
- $1083\ 00:52:17.490 \longrightarrow 00:52:19.380$ to find genes which correspond
- $1084\ 00:52:19.380$ --> 00:52:24.380 to this interaction between cancer and immune,
- $1085\ 00:52:25.100 --> 00:52:27.220$ and which are not necessarily markers of...
- 1086 00:52:27.220 --> 00:52:29.290 And regular markers of cancer and immune.
- 1087 00:52:29.290 --> 00:52:31.950 So, genes which are specifically related
- $1088\ 00:52:31.950 \longrightarrow 00:52:33.910$ to the non-linear interactions
- $1089\ 00:52:33.910 \longrightarrow 00:52:36.913$ between these patterns.
- 1090 00:52:37.810 --> 00:52:40.220 And to that end, basically we hypothesize
- 1091 00:52:40.220 --> 00:52:43.320 that since CoGAPS is already
- $1092\ 00:52:44.780 \longrightarrow 00:52:47.300$ an approximation of the dataset
- $1093\ 00:52:47.300 --> 00:52:49.720$ with a linear combination of the patterns,
- $1094\ 00:52:49.720 \longrightarrow 00:52:51.350$ the residuals of CoGAPS,
- $1095\ 00:52:51.350 --> 00:52:53.300$ of the CoGAPS estimate from the dataset

```
1096\ 00:52:54.810 --> 00:52:57.750 could point us to the non-linear interactions
```

- $1097\ 00:52:57.750 \longrightarrow 00:53:01.090$ between the patterns.
- $1098\ 00:53:01.090 \longrightarrow 00:53:04.950$ And we are only looking at the region
- $1099\ 00:53:06.410 --> 00:53:09.240$ where both of the patterns are active
- $1100\ 00:53:09.240 --> 00:53:11.753$ and comparing the residuals of CoGAPS
- $1101\ 00:53:12.870 \longrightarrow 00:53:15.330$ in that region to the residuals
- $1102\ 00:53:15.330 \longrightarrow 00:53:16.480$ in only the cancer region,
- $1103\ 00:53:16.480 \longrightarrow 00:53:18.350$ and only the immune region.
- $1104\ 00:53:18.350 \longrightarrow 00:53:21.293$ And now, this can be done for each of these,
- $1105\ 00:53:23.660 --> 00:53:24.840\ I$ guess, pattern combinations,
- $1106\ 00:53:24.840 \longrightarrow 00:53:28.810$ and we can find what corresponds
- $1107\ 00:53:28.810 \longrightarrow 00:53:30.010$ to pattern interaction
- $1108\ 00:53:30.010 \longrightarrow 00:53:32.340$ between these pairs of patterns.
- $1109\ 00:53:32.340 \longrightarrow 00:53:35.290$ So, for future work, as part
- 1110 00:53:35.290 --> 00:53:37.280 of the data collection in clinical trials,
- 1111 00:53:37.280 --> 00:53:42.280 we're already collecting both spacial
- $1112\ 00:53:42.380 \longrightarrow 00:53:44.440$ and single cell transcriptomics
- 1113 00:53:44.440 --> 00:53:47.260 and proteomics from patients.
- $1114\ 00:53:47.260 \longrightarrow 00:53:49.620$ So, we are trying to integrate all
- $1115\ 00:53:49.620 \longrightarrow 00:53:53.960$ of this into one big dataset,
- $1116\ 00{:}53{:}53.960 --> 00{:}53{:}56.673$ which would represent the tumor microenvironment.
- 1117 00:53:58.590 --> 00:54:00.550 which would help us characterize
- $1118\ 00:54:02.940 \longrightarrow 00:54:05.030$ the patient sample as a whole.
- 1119 00:54:05.030 --> 00:54:07.490 And we would also like
- $1120\ 00:54:07.490 \longrightarrow 00:54:10.470$ to infer intracellular signaling networks
- $1121\ 00:54:10.470 \longrightarrow 00:54:11.670$ the same way as we were trying to do
- 1122 00:54:11.670 --> 00:54:14.210 it using time, but now using space
- $1123\ 00:54:14.210 \longrightarrow 00:54:18.130$ where intracellular signaling is a function
- $1124\ 00:54:18.130 \longrightarrow 00:54:19.950$ of the distance between the cells
- $1125\ 00:54:19.950 \longrightarrow 00:54:21.410$ and the types of neighboring cells

- $1126\ 00:54:21.410 \longrightarrow 00:54:22.463$ for a target cell.
- $1127\ 00:54:25.549 \longrightarrow 00:54:27.130$ And the learnings from these projects
- 1128 00:54:27.130 --> 00:54:30.650 would go into a spatial temporal model
- 1129 00:54:30.650 --> 00:54:33.420 of tumor growth and response to therapy,
- $1130\ 00:54:33.420 \longrightarrow 00:54:35.880$ which can be used into building
- 1131 00:54:35.880 --> 00:54:39.370 a digital patient or digital clone,
- $1132\ 00:54:39.370 \longrightarrow 00:54:43.050$ where we can try to test what therapies
- $1133\ 00:54:43.050 \longrightarrow 00:54:47.093$ might work on what patients.
- $1134\ 00:54:48.430 \longrightarrow 00:54:50.350$ So, these are the people who have been,
- 1135 00:54:50.350 --> 00:54:51.540 and of course, 10x Genomics,
- $1136\ 00:54:51.540 \longrightarrow 00:54:53.950$ who were kind enough to give us the sample
- 1137 00:54:53.950 --> 00:54:58.510 for studying, as well as my collaborators
- $1138\ 00:54:58.510 \longrightarrow 00:54:59.593$ on this project.
- $1139\ 00:55:00.750 \longrightarrow 00:55:01.583$ Thank you so much.
- 1140 00:55:01.583 --> 00:55:03.010 And I can take questions now,
- $1141\ 00:55:04.350 \longrightarrow 00:55:05.950$ sorry for the overshooting time.
- $1142\ 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 00:55:09.776 --> 00:55:10.609 < v \text{ Lecturer} > \text{Thank you so much.} < /v > 1142 00:55:09.776 --> 0$
- $1143\ 00:55:10.609 \longrightarrow 00:55:15.320$ Do we have any questions to look at?
- 1144 00:55:22.275 --> 00:55:24.561 People on Zoom? Yeah, question (mumbles).
- 1145 00:55:24.561 --> 00:55:25.410 <v Female Student>Going back</v>
- $1146\ 00:55:25.410 \longrightarrow 00:55:27.676$ to the time series slides.
- 1147 00:55:27.676 --> 00:55:28.509 <v -> Mm-hmm. </v>
- 1148 00:55:28.509 --> 00:55:29.342 <v Female Student>Can you talk</v>
- 1149 00:55:29.342 --> 00:55:30.520 about how you know if you have good,
- $1150\ 00:55:30.520 \longrightarrow 00:55:32.240$ or bad pseudo times?
- $1151\ 00:55:32.240 \longrightarrow 00:55:35.040$ And is there a way to fix bad pseudo times?
- 1152 00:55:35.040 --> 00:55:38.820 <v ->So, yeah, as in what I've not shared on here </v>
- $1153\ 00:55:38.820 \longrightarrow 00:55:42.850$ is so, in our experiments,
- $1154\ 00:55:42.850 \longrightarrow 00:55:45.560$ we also, we knew for example,
- $1155\ 00:55:45.560 \longrightarrow 00:55:46.580$ that we were studying...

- 1156 00:55:46.580 --> 00:55:48.710 We wanted to study a trajectory which goes
- 1157 00:55:48.710 --> 00:55:53.710 from stem cells to neuroectoderm,
- $1158\ 00:55:55.710 \longrightarrow 00:55:57.120$ and we had markers.
- 1159 00:55:57.120 --> 00:55:59.713 And I think, some (mumbles) themselves.
- $1160\ 00:56:00.860 \longrightarrow 00:56:03.170$ They have identified markers
- 1161 00:56:03.170 \rightarrow 00:56:08.170 of stem cells neuroectoderms and endoderm cells.
- 1162 00:56:08.660 --> 00:56:10.270 So, if we're looking at the trajectories
- $1163\ 00:56:10.270 --> 00:56:13.040$ of the markers along the pseudo time
- $1164\ 00:56:13.040 \longrightarrow 00:56:14.493$ to see if those make sense.
- 1165 00:56:15.360 --> 00:56:17.800 For example, a marker which is supposed
- 1166 00:56:17.800 --> 00:56:21.250 to be high in stem cells would,
- 1167 00:56:21.250 --> 00:56:23.460 should be tapering down to zero
- 1168 00:56:23.460 --> 00:56:26.220 along pseudo time, and a marker,
- $1169\ 00:56:26.220$ --> 00:56:29.610 which is supposed to be high in neuroectoderm
- $1170\ 00:56:29.610 \longrightarrow 00:56:34.170$ should be increasing with pseudo time.
- 1171 00:56:34.170 --> 00:56:37.600 So, we had, I think six oral markers
- $1172\ 00:56:37.600 \longrightarrow 00:56:40.610$ to each of stem cells, neuroectoderm
- $1173\ 00:56:40.610 \longrightarrow 00:56:45.500$ and endoderm cells.
- $1174\ 00{:}56{:}45.500 \dashrightarrow 00{:}56{:}49.040$ And we were trying to confirm the combination
- $1175\ 00:56:49.040 --> 00:56:51.790$ that neuroectoderm markers increase
- $1176\ 00:56:51.790 --> 00:56:53.750$ with pseudo time, but the other two decrease,
- $1177\ 00:56:53.750 \longrightarrow 00:56:58.130$ or the endoderm shouldn't decrease necessarily,
- 1178 00:56:58.130 --> 00:57:00.300 but it shouldn't have
- $1179\ 00:57:00.300$ --> 00:57:05.300 a monotonic increase like the neuroectoderm one.
- 1180 00:57:08.280 --> 00:57:10.903 And it should not be present in the initial.
- 1181 00:57:12.380 --> 00:57:13.213 Does that...
- $1182\ 00:57:14.310 \longrightarrow 00:57:16.360$ So, that was one way to do it, basically.

```
1183 00:57:21.120 --> 00:57:22.116 <v Lecturer>Thank you.</v>
```

- 1184 00:57:22.116 --> 00:57:23.783 Any other questions?
- 1185 00:57:39.510 --> 00:57:41.423 So, with the combination of many cells,
- $1186\ 00:57:41.423 \longrightarrow 00:57:43.630$ and the spatial stuff, is there any hope
- $1187\ 00:57:43.630 \longrightarrow 00:57:45.740$ of getting a temporal signal out of any of that,
- $1188\ 00:57:45.740 \longrightarrow 00:57:46.940$ or is that (indistinct)?
- $1189\ 00:57:49.897 --> 00:57:52.060 < v -> In spatial did you mean? </v>$
- 1190 00:57:52.060 --> 00:57:53.350 <v Lecturer>Yeah.</v>
- 1191 00:57:53.350 --> 00:57:58.350 <v Dr. Deshpande>So, I think,</v>
- 1192 00:57:58.840 --> 00:58:00.060 the issue would be, I guess,
- $1193\ 00:58:00.060 \longrightarrow 00:58:01.653$ not in clinical, I suppose.
- 1194 00:58:03.740 --> 00:58:05.740 In a sense that, okay, are you thinking
- 1195 00:58:05.740 --> 00:58:08.045 about pseudo temporal, or just clinical?
- 1196~00:58:08.045 --> 00:58:09.700 < v Lecturer>Yeah.</v>
- 1197 00:58:09.700 --> 00:58:11.330 <-> Pseudo temporal, I think there might <-/> /v>
- 1198 00:58:11.330 --> 00:58:12.290 be some possibility,
- 1199 00:58:12.290 --> 00:58:13.570 and I've been thinking of
- 1200 00:58:17.940 --> 00:58:19.500 as in, we would still have to isolate,
- $1201\ 00:58:19.500 \longrightarrow 00:58:21.610\ I$ guess, cell types, for example.
- $1202\ 00:58:21.610 \longrightarrow 00:58:24.130$ So, one of the problems with that
- $1203\ 00:58:24.130 \longrightarrow 00:58:26.640$ is that as I mentioned,
- 1204 00:58:26.640 --> 00:58:30.550 the spots are not exactly single cell, right?
- 1205 00:58:30.550 --> 00:58:32.800 So, especially, let's say if you're trying
- $1206\ 00:58:32.800 --> 00:58:35.500$ to do a pseudo temporal ordering
- 1207 00:58:35.500 --> 00:58:38.283 of CD8 T-cells,
- $1208\ 00:58:39.380 \longrightarrow 00:58:40.500$ they are more,
- $1209\ 00:58:40.500 \longrightarrow 00:58:44.460$ more likely than not, co-localized
- 1210 00:58:44.460 --> 00:58:48.830 with other cell types, which would also,
- 1211 00:58:48.830 --> 00:58:51.250 I guess, corrupt the expression

- $1212\ 00:58:51.250 \longrightarrow 00:58:53.030$ that you are seeing.
- 1213 00:58:53.030 --> 00:58:57.180 So, that would make it slightly different.
- 1214 00:58:57.180 --> 00:59:01.300 We could think of ordering the spot
- $1215\ 00:59:01.300 \longrightarrow 00:59:02.944$ as a whole, basically.
- 1216 00:59:02.944 --> 00:59:03.777 And my...
- 1217 00:59:04.910 --> 00:59:07.120 I belong to a school of thought that basically,
- 1218 00:59:07.120 --> 00:59:08.446 if you have a...
- 1219 00:59:08.446 --> 00:59:09.600 And then, so what people try to do
- 1220 00:59:09.600 --> 00:59:13.000 with say, this kind of data,
- 1221 00:59:13.000 --> 00:59:14.560 this spacial Visium data,
- 1222 00:59:14.560 --> 00:59:16.830 where you have say, up to 10 cells,
- $1223\ 00:59:16.830 \dashrightarrow 00:59:21.830$ they try to resolve this into cell types.
- $1224\ 00:59:22.730 \longrightarrow 00:59:25.010$ So, they would compare that to, there is I think,
- $1225\ 00:59:25.010 \longrightarrow 00:59:27.703$ one paper called RTCD, or RCTD.
- 1226 00:59:29.940 --> 00:59:32.223 RCTD robust cell type decomposition.
- $1227\ 00:59:33.470 \longrightarrow 00:59:35.510$ So, what they do is basically,
- $1228\ 00:59:35.510 \longrightarrow 00:59:37.870$ they take the spatial data,
- 1229 00:59:37.870 --> 00:59:41.060 they have a reference single cell data,
- $1230\ 00:59:41.060 \longrightarrow 00:59:46.060$ and they try to assign each spot,
- $1231\ 00:59:46.990 --> 00:59:51.030$ or a resolve each spot into a mixture
- 1232 00:59:51.030 --> 00:59:53.930 of the cell types that might exist
- $1233\ 00:59:53.930 \longrightarrow 00:59:55.543$ in the single cell data.
- 1234 00:59:56.750 --> 01:00:01.010 And that could help you to say,
- $1235\ 01:00:01.010 --> 01:00:04.380$ identify what the mixture in general is.
- $1236\ 01:00:04.380 \longrightarrow 01:00:08.960$ But my as in my thought is that we could
- 1237 01:00:08.960 --> 01:00:12.970 just think of each spot as some representation
- $1238\ 01:00:15.200 \longrightarrow 01:00:16.820$ of the biology in that neighborhood.
- 1239 01:00:16.820 --> 01:00:19.710 So, each spot could just represent
- 1240 01:00:19.710 --> 01:00:22.360 a neighborhood, as opposed to trying to find
- $1241\ 01:00:22.360 \longrightarrow 01:00:23.893$ what the individual cells are.

- 1242 01:00:24.990 --> 01:00:28.840 And that would basically abstract out
- 1243 01:00:30.460 --> 01:00:33.340 the representation and the biology to that
- $1244\ 01:00:33.340 \longrightarrow 01:00:34.900$ of the spots.
- $1245\ 01:00:34.900 \longrightarrow 01:00:36.790$ And we'll have to think about how to do that,
- $1246\ 01:00:36.790 --> 01:00:40.350$ but I think there could be some ordering to that,
- $1247\ 01:00:40.350 \longrightarrow 01:00:45.130$ but we'll need to see what makes sense.
- 1248 01:00:45.130 --> 01:00:49.410 And then, for a lot of cells, cell states,
- $1249\ 01:00:49.410 \longrightarrow 01:00:50.650$ they are quite well-characterized.
- 1250 01:00:50.650 --> 01:00:52.670 For example, if you say that a T-cell
- 1251 01:00:52.670 --> 01:00:55.210 is activated, or a T-cell as naive,
- $1252\ 01:00:55.210$ --> 01:00:59.150 or exhausted, you know what markers to expect.
- $1253\ 01:00:59.150 \longrightarrow 01:01:01.400$ But what would you be able to say
- $1254\ 01:01:02.340 \longrightarrow 01:01:04.043$ for spots instead?
- 1255 01:01:05.420 --> 01:01:09.900 The other thing to think of is,
- 1256 01:01:09.900 --> 01:01:12.320 especially with say, the proteomics as well,
- 1257 01:01:12.320 --> 01:01:14.370 where you can get actual single cell
- $1258\ 01:01:18.000 --> 01:01:22.380$ and distributions, and neighborhood characterization.
- 1259 01:01:22.380 --> 01:01:25.033 You could think of it as can you,
- $1260\ 01:01:26.810 \longrightarrow 01:01:28.347$ so the same thing that...
- $1261\ 01:01:28.347 \longrightarrow 01:01:30.620$ The same ideas that were used
- 1262 01:01:30.620 --> 01:01:32.743 for pseudo temporal ordering of cells,
- $1263\ 01:01:33.590 --> 01:01:35.720$ can they be used for pseudo temporal
- 1264 01:01:35.720 --> 01:01:38.680 ordering of neighborhoods?
- 1265 01:01:38.680 --> 01:01:40.730 For example, if you have a cell neighborhood,
- 1266 01:01:40.730 --> 01:01:44.868 which as they're presented as whatever,
- 1267 01:01:44.868 --> 01:01:47.763 the central cell, and it's five neighbors.
- 1268 01:01:48.710 --> 01:01:51.540 Now, depending on, are they all tumor?
- $1269\ 01:01:51.540 \longrightarrow 01:01:52.860$ Then maybe they have...

- 1270 01:01:52.860 --> 01:01:54.310 They're basically deep in the cancer,
- $1271\ 01:01:54.310 --> 01:01:57.183$ which has never been visited by an immune cell,
- 1272 01:01:58.140 --> 01:01:59.380 is that a mix of tumor
- 1273 01:01:59.380 --> 01:02:01.570 and activated immune cells?
- 1274 01:02:01.570 --> 01:02:03.930 So, that is basically an active tumor
- 1275 01:02:03.930 --> 01:02:06.150 immune interaction that's happening.
- 1276 01:02:06.150 --> 01:02:10.220 Is that exhausted T-cells and tumor,
- $1277\ 01:02:10.220 \longrightarrow 01:02:11.100$ where basically the tumor
- 1278 01:02:11.100 --> 01:02:15.690 has fought back and tried to suppress the...
- 1279 01:02:15.690 --> 01:02:16.930 Or it's basically sent signals
- $1280\ 01:02:16.930 \longrightarrow 01:02:21.130$ to suppress the immune response, and so on.
- 1281 01:02:21.130 --> 01:02:22.433 So, perhaps there could be
- 1282 01:02:22.433 --> 01:02:24.730 a trajectory of neighborhoods,
- 1283 01:02:24.730 --> 01:02:28.810 where you could say that depending on all
- 1284 01:02:28.810 --> 01:02:31.130 the possible combinations that you expect
- 1285 01:02:31.130 --> 01:02:33.453 in cellular neighborhoods,
- 1286 01:02:35.330 --> 01:02:39.960 this current neighborhood is this far along
- 1287 01:02:39.960 --> 01:02:42.680 that process, or that branch of a process.
- $1288\ 01:02:44.393 --> 01:02:46.621$ That was a long and winding answer.
- $1289\ 01:02:46.621 \longrightarrow 01:02:47.740$ (chuckles) I don't know if
- 1290 01:02:48.680 --> 01:02:51.690 that necessarily answered it. <v Lecturer>Thank you.</v>
- 1291 01:02:51.690 --> 01:02:54.490 Thank you, any last questions?
- $1292\ 01:02:54.490 --> 01:02:55.770\ I$ wanna be mindful of time.
- 1293 01:02:55.770 --> 01:02:58.333 Any questions that come to you, or?
- 1294 01:03:06.287 --> 01:03:09.277 All right, well if not, thank you again.
- 1295 01:03:09.277 --> 01:03:11.168 (students applaud) We really appreciate that.
- $1296\ 01:03:11.168 --> 01:03:14.752 < v$ Dr. Deshpande>Thank you a lot.</v>
- 1297 01:03:14.752 --> 01:03:15.977 <v Lecturer>You have a wonderful (indistinct).</v>
- 1298 01:03:15.977 --> 01:03:16.810 <v -> Mm-hmm. </v>

 $1299\ 01:03:20.394 \dashrightarrow 01:03:24.394\ (lecturer\ mumbles\ indistinctly)$ $1300\ 01:03:26.984 \dashrightarrow 01:03:31.067\ (students\ chatter\ indistinctly)$